

**TECHNICAL SHEET**

<b>AS006</b>	<b>MIU Medium Slant</b>				
<b>Formula</b>					
<b>Ingredients:</b>		<b>gms/lit.</b>			
Casein enzymic hydrolysate		10.00			
Dextrose		1.00			
Sodium chloride		5.00			
Phenol red		0.01			
Agar		2.00			
Final pH (at 25°C): 6.8 ± 0.2					
<b>Directions:</b>					
Streak the test inoculum aseptically into the slant and incubate at appropriate conditions.					
<b>Principle:</b>					
MIU Medium Base is formulated to detect motility, urease and indole production in single tube. Casein Enzymic Hydrolysate provides amino acids and other nitrogenous substance. Dextrose is the fermentable carbohydrate. Sodium chloride maintains osmotic equilibrium. Phenol red is the pH indicator. When urea is utilized, ammonia is formed during incubation which makes the medium alkaline showing a pink - red color by the change in phenol red indicator. Motility and urease reactions are read before testing Indole production. Indole is produced from tryptophan present in casein enzymic hydrolysate. The Indole produced combines with the aldehyde present in the Kovac's reagent to form a red complex. Motile organisms show either diffused growth or turbidity away from stab line.					
<b>(I) QC Tests</b>					
	pH:	6.8 ± 0.2			
	Color:	Orange coloured opalescent butts			
	Appearance:	Sterile MIU Medium in disposable slants.			
<b>(II) Sterility test</b>					
Passes release criteria					
<b>(III) Q.C. Test Microbiological</b>					
Cultural characteristics observed after incubation at 35-37°C for 18-24 hours.					
	MICROORGANISM (ATCC)	GROWTH	MOTILITY	INDOLE	UREASE
	ESCHERICHIA COLI 25922	Luxuriant	Positive	Positive	Negative
	Klebsiella pneumoniae 13883	Luxuriant	Negative	Negative	Weakly positive
	Proteus vulgaris 13315	Luxuriant	Positive	Positive	Positive
	Proteus mirabilis 25933	Luxuriant	Positive	Negative	Positive
	S. serotype typhimurium 14028	Luxuriant	Positive	Negative	Negative

Refer disclaimer Overleaf

**TECHNICAL SHEET**

<b>Precautions :</b>	1. In Vitro diagnostic use only. 2. Read the label before opening the container
<b>Limitations :</b>	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.
<b>Use:</b>	For detection of motility, urease and indole production.
<b>Storage:</b>	Store between 2-8°C. Use before expiry date on the label.
<b>Packing:</b>	10/25 disposable slants.

**Disclaimer:**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.