

TECHNICAL SHEET

B035	PEPTONE WATER				
Formula					
Ingredients:		gms/lit.			
Peptic digest of animal tissue	10.00				
Sodium chloride	5.00				
Final pH (at 25°C) : 7.2 ± 0.2					
Directions :					
Suspend 15 grams in 1000 ml. distilled water. Add the test carbohydrate in desired quantity and dissolve completely. Dispense in tubes with or without inverted Durham's tubes and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.					
Principle :					
Peptic digest of animal tissue provides essential nutrients. To study the fermentation ability of carbohydrates saccharose, rhamnose, salicin is generally added in 0.5% amount separately to the basal medium before or after sterilization. Sodium chloride maintains osmotic equilibrium.					
QC Tests – (I)Dehydrated Medium					
	Colour :	Cream to light yellow			
	Appearance :	Homogeneous Free Flowing powder			
(II)Rehydrated medium					
	pH (post autoclaving/heating) :	7.2 ± 0.2			
	Colour (post autoclaving/heating) :	Light amber to light yellow			
	Clarity (post autoclaving/heating) :	Clear to very slightly opalescent			
(III)Q.C. Test Microbiological					
	Cultural characteristics observed after 18 – 24 hrs.at 35 – 37°C.				
	MICROORGANISM (ATCC)	GROWTH	INDOLE TEST		
	Escherichia coli (25922)	Luxuriant	positive reaction, red ring at the interface of the medium on addition of Kovac's reagent (BA068)		
	Salmonella typhimurium (14028)	Luxuriant	negative reaction, no red ring at the interface of the medium on addition of Kovac's reagent (BA068)		
	Staphylococcus aureus (25923)	Luxuriant	negative reaction, no red ring at the interface of the medium on addition of Kovac's reagent (BA068)		
Precautions :	1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use:	As all-purpose growth medium and as the basis of carbohydrate fermentation media.				
Storage :	Dehydrated medium- below 30°C Prepared medium– Between 2 to 8°C.				
Packing :	500 gm. bottle				
Product profile:	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B035	15 g/l	33.333L	7.2 ± 0.2	NIL	121 ⁰ C / 15 minutes

BIOMARK Laboratories-INDIA

www.biomarklabs.com

TECHNICAL SHEET

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.