

BIOMARK Laboratories-INDIA

www.biomarklabs.com

TECHNICAL SHEET

B037	Potato Dextrose Broth				
Formula					
Ingredients :		gms/lit.			
Potato infusion from		200.00			
Dextrose		20.00			
Final pH (at 25°C) : 5.1 ± 0.2					
Directions :					
Suspend 24 grams in 1000 ml distilled water. Heat if necessary, to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well before dispensing. In specific work, when pH 3.5 is required, acidify the medium with sterile 10% tartaric acid. The amount of acid required for 100 ml. of sterile, cooled medium is approximately 1 ml. Do not heat the medium after addition of the acid.					
Principle :					
Potato infusion and dextrose promote luxuriant fungal growth. Adjusting the pH of the medium by tartaric acid to 3.5, inhibits the bacterial growth. Heating the medium after acidification should be avoided.					
QC Tests – (I) Dehydrated Medium					
	Colour :	Off-white to yellow			
	Appearance :	Homogeneous Free Flowing powder			
(II) Rehydrated medium					
	pH (post autoclaving/heating) :	5.1 ± 0.2			
	Colour (post autoclaving/heating) :	Light amber			
	Clarity (post autoclaving/heating) :	Clear to slightly opalescent			
(III) Q.C. Test Microbiological					
Cultural characteristics observed after an incubation at 25-30°C for 4-5 days					
	MICROORGANISM (ATCC)	GROWTH			
	Aspergillus niger (16404)	Luxuriant			
	Candida albicans (10231)	Luxuriant			
	Saccharomyces cerevisiae (9763)	Luxuriant			
Precautions :	1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :	For isolation and enumeration of yeasts and molds from dairy and other food products.				
Storage :	Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
Packing :	500 gm. bottle				
Product profile:	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B037	24g/l	20.833L	5.1 ± 0.2	NIL	121°C / 15 minutes

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARK LABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.