

BIOMARK Laboratories-INDIA

www.biomarklabs.com

TECHNICAL SHEET

B069	POTATO DEXTROSE AGAR				
Formula					
Ingredients :		gms/lit.			
Potatoes infusion from		300.00			
Dextrose		20.00			
Agar		15.00			
Final pH (at 25°C) :		5.6 ± 0.2			
Directions :					
Suspend 41 grams in 1000 ml. distilled water. Warm gently to dissolve medium completely. Dispense in tubes or flasks. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well before dispensing.					
Principle :					
Potato infusion and dextrose promote luxuriant fungal growth. Adjusting the pH of the medium by tartaric acid inhibits the bacterial growth. Heating the medium after acidification should be avoided as it may hydrolyse the agar which can render the agar unable to solidify.					
QC Tests – (I)Dehydrated Medium					
	Colour :	Cream to light yellow			
	Appearance :	Homogeneous Free Flowing powder			
(II)Rehydrated medium					
	pH (post autoclaving/heating) :	5.6 ± 0.2			
	Colour (post autoclaving/heating) :	Light amber			
	Clarity (post autoclaving/heating) :	Clear to slightly opalescent			
(III)Q.C. Test Microbiological					
	Cultural characteristics observed after 4 – 5 days at 22 - 25°C.				
	MICROORGANISM (ATCC)	GROWTH			
	Aspergillus niger (16404)	Luxuriant			
	Candida albicans (10231)	Luxuriant			
	Saccharomyces cerevisiae (9763)	Luxuriant			
Precautions :	1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium. 2. Heating Potato Dextrose Agar after acidifying hydrolyzes the agar and may destroy the solidifying properties. 3. Potato Dextrose Agar is not a differential medium. Perform microscopic examination and biochemical tests to identify isolates to genus and species if necessary.				
Use :	For cultivation and enumeration of yeasts and molds from dairy and other food products.				
Storage :	Dehydrated medium- below 30°C Prepared medium– Between 2 to 8°C.				
Packing :	500 gm. bottle				
Product profile:	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B069	41g/l	12.195L	5.6 ± 0.2	NIL	121°C / 15 minutes

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.