

**TECHNICAL SHEET**

<b>B085</b>	<b>TELLURITE BLOOD AGAR BASE</b>				
<b>Formula</b>					
<b>Ingredients :</b>		<b>gms/lit.</b>			
Biopeptone					10.00
Sodium chloride					5.00
Dipotassium hydrogen phosphate					4.00
Corn starch					1.00
Monopotassium phosphate					1.00
Agar					10.00
Final pH (at 25°C) : 7.2 ± 0.2					
<b>Directions :</b>					
Suspend 31 grams in 500 ml distilled water to make a double strength base. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Aseptically add sterile prepared Haemoglobin solution (BF036) (10 grams in 490ml distilled water) and sterile reconstituted contents of one vial of Vitamino Growth Supplement (BF037) and 1%Potassium Tellurite (BF010). Mix well and pour into sterile Petri plates.					
<b>Principle :</b>					
Biopeptone provides nitrogenous compounds. Sodium chloride maintains the osmotic equilibrium of the medium while phosphates buffer the medium. Corn starch neutralizes the toxic metabolites. Haemoglobin and Vitamino Growth Supplement stimulate good growth of Corynebacterium. Potassium tellurite acts as a selective agent and has inhibitory activity against most gram-positive and gram-negative bacteria except Corynebacterium species. C.diphtheriae reduces potassium tellurite to tellurium and thereby produce gray-black coloured colonies.					
<b>QC Tests - (I)Dehydrated Medium</b>					
	Colour :	Cream to yellow			
	Appearance :	Homogeneous Free Flowing powder			
<b>(II)Rehydrated medium</b>					
	pH (post autoclaving/heating) :	7.2 ± 0.2			
	Colour (post autoclaving/heating) :	a) Yellow b) With the addition of haemoglobin : Reddish brown			
	Clarity (post autoclaving/heating) :	a) Clear b) Opaque			
<b>(III)Q.C. Test Microbiological</b>					
	Cultural characteristics observed along with added supplements after an incubation at 35-37°C for 48 hours (or more).				
	MICROORGANISM (ATCC )	GROWTH	COLOUR OF COLONY		
	Corynebacterium diphtheriae (11913)	Good – luxuriant	Grey – black		
	Escherichia coli (25922)	Inhibited	-		
<b>Precautions :</b>	1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
<b>Limitations :</b>	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
<b>Use :</b>	For the selective isolation and cultivation of Corynebacterium species.				
<b>Storage :</b>	Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
<b>Packing :</b>	500 gm. bottle				
<b>Product profile:</b>	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
<b>B085</b>	31g/l	16.129L	7.2 ± 0.2	Haemoglobin (BF036) Vitamino Growth Supplement (BF037) 1% Potassium Tellurite (BF010).	121°C /15 min.

Refer disclaimer Overleaf

**Disclaimer:**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.