

TECHNICAL SHEET

B1009	DICHLORAN GLYCEROL MEDIUM BASE				
Formula					
Ingredients :		gms/lit.			
Peptic digest of animal tissue	5.00				
Dextrose		10.00			
Monopotassium phosphate		1.00			
Magnesium sulphate		0.50			
Chloramphenicol		0.10			
Dichloran		0.002			
Agar		15.00			
Final pH (at 25°C) : 5.6 ± 0.2					
Directions :					
Suspend 15.8 grams in 500 ml distilled water. Heat to boiling to dissolve the medium completely. Add 110 grams of glycerol (Analytical Reagent Grade). Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour into sterile Petri plates.					
Principle :					
Dichloran Glycerol Medium is recommended for isolation and enumeration of xerophilic moulds from dried and semidried foods. Peptic digest of animal tissue provides nitrogen, vitamins and minerals. Dextrose is a carbohydrate source. Phosphate buffers the medium. Magnesium sulfate provides divalent cations and sulfate. Dichloran is an antifungal agent, added to the medium to reduce colony diameters of spreading fungi. The glycerol at 18% (w/w) lowers the water activity Chloramphenicol inhibits gram – negative and gram – positive bacteria.					
QC Tests – (I) Dehydrated Medium					
Colour :	Cream to light yellow				
Appearance :	Homogeneous Free Flowing powder				
(II) Rehydrated medium					
PH (post autoclaving/heating) :	5.6 ± 0.2				
Colour (post autoclaving/heating) :	Medium amber				
Clarity (post autoclaving/heating) :	Slightly opalescent				
(III) Q.C. Test Microbiological					
Cultural characteristics observed with added 22 grams of glycerol after an incubation at 25°C for upto 6 days.					
MICROORGANISM (ATCC)	GROWTH				
Mucor racemosus (42647)	Good - luxuriant				
Saccharomyces cerevisiae (9763)	Good – luxuriant				
Candida albicans (10231)	Good – luxuriant				
Escherichia coli (25922)	Inhibited				
Bacillus subtilis (6633)	Inhibited				
Precautions :	1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :	For selective isolation of xerophilic molds from food samples.				
Storage :	Dehydrated medium- below 30°C Prepared medium– Between 2 to 8°C.				
Packing :	500 gm bottle				
Product profile:	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B1009	31.6g/l	15.82L	5.6 ± 0.2	Glycerol	121°C / 15 minutes

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARK LABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.