

BIOMARK Laboratories-INDIA

www.biomarklabs.com

TECHNICAL SHEET

B1056	HALOPHILIC BROTH					
Formula						
Ingredients :		gms/lit.				
Casein acid hydrochloride		10.00				
Yeast extract		10.00				
Proteose peptone		5.00				
Trisodium citrate		3.00				
Potassium chloride		2.00				
Magnesium sulphate		25.00				
Sodium chloride		250.00				
Final pH (at 25°C) : 7.2 ± 0.2						
Directions :						
Suspend 30.5 gms in 100 ml distilled water. Heat to boiling to dissolve the medium completely. Dispense as desired and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.						
Principle :						
Halophilic media contains casein acid hydrolysate, proteose peptone and yeast extract which provide all the necessary nutrients, mainly nitrogenous and vitamins to the halophilic bacteria. Trisodium citrate is added to avoid the losses. Magnesium is an essential ion for the growth of extreme halophiles and is incorporated in the medium as magnesium sulphate.						
QC Tests – (I) Dehydrated Medium						
Colour :		Light yellow				
Appearance :		Homogeneous Free Flowing powder				
(II) Rehydrated medium						
pH (post autoclaving/heating) :		7.2 ± 0.2				
Colour (post autoclaving/heating) :		Amber				
Clarity (post autoclaving/heating) :		Slightly opalescent gel/hazy solution				
(III) Q.C. Test Microbiological						
Cultural characteristics observed after 12 days at 35-37°C.						
MICROORGANISM (ATCC)		GROWTH				
Halococcus morrhuae (17082)		Luxuriant				
Halobacterium salinarium (33171)		Luxuriant				
Precautions :		1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :		For isolation and cultivation of extremely halophilic bacteria.				
Storage :		Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
Packing :		500 gm bottle				
Product profile:		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B1056		305.0 g/l	1.639 L	7.2± 0.2	Nil	121°C/15min.

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARK LABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.