

TECHNICAL SHEET

B1060	HERELLEA AGAR					
Formula						
Ingredients :		gms/lit.				
Casein enzymic hydrolysate		15.00				
Papaic digest of soyabean meal	5.00					
Sodium chloride		5.00				
Lactose		10.00				
Maltose		10.00				
Bile salt mixture		1.25				
Bromo cresol purple		0.02				
Agar		16.00				
Final pH (at 25°C) : 6.8 ± 0.2						
Directions :						
Suspend 62.27 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.						
Principle :						
Casein enzymic hydrolysate and papaic digest of soyabean meal are sources of carbon, nitrogen, vitamins and minerals. Sodium chloride provides the essential ions and also maintains the osmotic equilibrium of the medium. Bile salts mixture in the medium acts as selective agent, inhibiting the growth of Neisseria species and other gram-positive organisms. Lactose and maltose are the fermentable carbohydrates. Bromocresol purple acts as the pH indicator.. Fermentative gram-negative bacteria ferment the carbohydrates to produce acid, which cause a corresponding change in the colour of pH indicator dye to yellow.						
QC Tests - (I)Dehydrated Medium						
Colour :		Cream to yellow				
Appearance :		Homogeneous Free Flowing powder				
(II)Rehydrated medium						
pH (post autoclaving/heating) :		6.8 ± 0.2				
Colour (post autoclaving/heating) :		Purple				
Clarity (post autoclaving/heating) :		Clear to slightly opalescent				
(III)Q.C. Test Microbiological						
Cultural characteristics observed after 18 -24 hoursat 35-37°C.						
MICROORGANISM (ATCC)		GROWTH	COLOUR			
Acinetobacter calcoaceticus (17961)		Luxuriant	Pale lavender			
Acinetobacter lwoffii (9957)		Luxuriant	Pale lavender			
Escherichia coli (25922)		Luxuriant	Yellow			
Listeria monocytogenes (19112)		Inhibited	-			
Staphylococcus aureus (25923)		Inhibited	-			
Precautions :		1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :		For the selective isolation and differentiation of gram-negative, fermentation and non-fermentative organisms especially for differentiation of organisms of Mima and Herellea group.				
Storage :		Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
Packing :		500 gm bottle				
Product profile:		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B1060		62.27 g/l	16.05 L	6.8± 0.2	Nil	121°C/15min.

Refer disclaimer Overleaf

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.