BIOMARK Laboratories-INDIA

www.biomarklabs.com

TECHNICAL SHEET

B1070 INHIBITORY MOLD AGAR, ULRICH (MOLD INHIBITORY AGAR, ULRICH)							
Formula							
Ingredients: gms/lit. Casein enzymic hydrolysate 3.00							
			3.00 2.00				
Peptic digest of animal tissue Yeast extract			5.00				
Dextrose			5.00				
Starch, soluble			2.00				
Dextrin			1.0				
Sodium phosphate			2.0				
Ferrous sulphate			0.04				
Magnesium sulphate			0.8				
Sodium chloride			0.04				
Manganese sulphate			0.16				
Chloramphenicol			0.125				
Agar			15.	00			
Final pH (at 25°C) :6.7±0.2							
Directions:							
Suspend 36.17 grams in 1000 ml distilled water. Mix thoroughly and heat to boiling to dissolve the							
medium completely. Sterilize by autoclaving at 118 - 121°C (12-15 lbs pressure) for 15 minutes. Mix well							
and pour into sterile Petri plates.							
Principle:							
Casein enzymic hydrolysate and peptic digest of animal tissue provide essential growth nutrients. Yeast							
extract is a rich source of vitamin B complex. Dextrose, starch and dextrin are energy sources for the							
metabolism of fungi. Sodium chloride and metallic salts provide essential ions and minerals.							
Chloramphenicol inhibits a wide variety of gram-positive and gram-negative bacteria. Potential							
contaminants of cosmetics and toiletries like Pseudomonas aeruginosa and Serratia marcescens are							
effectively inhibited by chloramphenicol. Sodium phosphates buffer the medium.							
QC Tests - (I)Dehydrated Medium							
Colour :			Cream to yellow				
Appearance :			Homogeneous Free Flowing powder				
(II)Rehydrated medium							
pH (post autoclaving/heating) :			6.7±0.2				
Colour (post autoclaving/heating):			Amber				
Clarity (post autoclaving/heating):			Clearto slightly opalescent gel				
(III)Q.C. Test Microbiological							
Cultural characteristics observed after an incubation at 25-30°C for upto 7 days ii)Bacterial cultures are							
incubated at 35-37°C							
			ROWTH				
` '			xuriant				
` ′			hibited				
	hibited						
Trichophyton mentagrophytes (9533) luxuriant							
Precautions :							
	2. Follow proper, established laboratory procedures in handling and disposing of infection						
materials. Limitations: 1. Since the nutritional requirements of organisms vary, some strains may be							
Limitations : 1. Since the nutritional requirements of organisms vary, some strains ma encountered that fail to grow or grow poorly on this medium.						ailis Illay De	
Use :	It is used for selective isolation of pathogenic fungi.						
Storage: Dehydrated medium- at 15-25°C Prepared medium- Between 2 to 8°C.						0 8°C	
Packing: 500 gm bottle							
Product			n	рН	Supplement	Sterilization	
		Preparatio		(25°C)	Заррістісті	Stermzation	
B1070	36.17 g/l	13.823 L		6.7±0.2	Nil	118-121°C/15 minutes	
	33.17 9/1	15.0		3.7 = 3.2			

Refer disclaimer Overleaf page 01 of 02

BIOMARK Laboratories-INDIA www.biomarklabs.com

TECHNICAL SHEET

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

Page 02 of 02

Rev: July 2021