

TECHNICAL SHEET

B1083	MYCOLOGICAL AGAR, MODIFIED					
Formula						
Ingredients :		gms/lit.				
Papain digest of soyabean meal 10.00						
Dextrose		10.00				
Agar		16.00				
Final pH (at 25°C) :		7.0 ± 0.2				
Directions :						
Suspend 36 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour into sterile Petri plates.						
Principle :						
Papain digest of soyabean meal serves as source of carbon, nitrogen and other essential growth nutrients. Dextrose is the source of energy. Agar is the solidifying agent. The pH may be adjusted to 4.0 after autoclaving by adding sterile 10% lactic acid/acetic acid and used for determining yeast and mold counts of carbonated beverages and food products						
QC Tests – (I) Dehydrated Medium						
Colour :		Cream to yellow				
Appearance :		Homogeneous Free Flowing powder				
(II) Rehydrated medium						
pH (post autoclaving/heating) :		7.0 ± 0.2				
Colour (post autoclaving/heating) :		Light amber				
Clarity (post autoclaving/heating) :		Clear to slightly opalescent				
(III) Q.C. Test Microbiological						
Cultural characteristics observed after 48- 72 hrs.at 25-30°C.						
MICROORGANISM (ATCC)		Growth on Agar w/ low pH		Growth		
Aspergillus niger (16404)		Good				
Candida albicans (10231)		Good		Good		
Lactobacillus acidophilus (11506)		Good		Good		
Staphylococcus aureus (25923)		Inhibited		Good		
Saccharomyces cerevisiae (9763)		Good		Good		
Precautions :		1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :		It is used for cultivation of fungi.				
Storage :		Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
Packing :		500 gm. bottle				
Product profile:		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B1083		36g/l	13.888 L	7.0 ± 0.2	NIL	121°C / 15 minutes

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.