

TECHNICAL SHEET

B1159	CAMPYLO THIOGLYCOLLATE MEDIUM				
Formula					
Ingredients :		gms/lit.			
Casein enzymic hydrolysate		20.00			
Sodium chloride		2.50			
Dipotassium phosphate	1.50				
Sodium thioglycollate	0.60				
L-Cystine		0.40			
Sodium sulphite		0.20			
Agar		1.60			
Final pH (at 25°C) : 7.0 ± 0.2					
Directions :					
Suspend 26.8 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. To make the medium selective for Campylobacter species, add reconstituted contents of 2 vials of Campylobacter supplement-I (Blaser-Wang, BF013). Mix well before dispensing.					
Principle :					
The medium contains necessary nutrients to promote growth of Campylobacter species. Moreover the supplement (Blaser and Wang) consists of five antibiotics viz. Amphotericin B, Cephalothin, Polymyxin B, Trimethoprim and Vancomycin which inhibit multiplication of normal microbial flora in faecal specimens thus facilitating isolation of Campylobacter jejuni, Cephalothin may not always inhibit Campylobacter fetus species and some strains may grow at 42°C. Further tests should be performed to confirm the Campylobacter jejuni.					
QC Tests – (I)Dehydrated Medium					
Colour :		Yellow			
Appearance :		Homogeneous Free Flowing powder			
(II)Rehydrated medium					
pH (post autoclaving/heating) :		7.0 ± 0.2			
Colour (post autoclaving/heating) :		Light to medium amber			
Clarity (post autoclaving/heating) :		Very slightly opalescent			
(III)Q.C. Test Microbiological					
Cultural characteristics observed with added Campylobacter Supplement I(Blaser Wang, BF013) in an atmosphere of 5-15% O2 and 5-12% CO2 after an incubation at 42°C for 18-24 hours.					
MICROORGANISM (ATCC)		GROWTH			
Campylobacter jejuni (33291)		Good-luxuriant			
Campylobacter coli (33559)		Good-luxuriant			
Helicobacter pylori (43504)		Good-luxuriant			
Escherichia coli (25922)		None – poor			
Precautions :	1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use:	For maintenance, transport and storage of Campylobacter species				
Storage :	Dehydrated medium- below 30°C Prepared medium– Between 2 to 8°C.				
Packing :	500 gm bottle				
Product profile:	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B1159	26.8g/l	18.65L	7.0 ± 0.2	Campylobacter supplementI(Blaser Wang, BF013)	121°C / 15 minutes

Refer disclaimer overleaf

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.