

**TECHNICAL SHEET**

<b>B1201</b>	<b>LEES MULTIDIFFERENTIAL AGAR</b>				
<b>Formula</b>					
<b>Ingredients :</b>		<b>gms/lit.</b>			
Tomato Juice broth		41.00			
Peptonized milk		20.00			
Calcium pantothenate		2.00			
Citric acid		1.10			
Calcium carbonate		5.00			
Polysorbate 80		0.50			
Bromo cresol green		0.022			
Cycloheximide		0.007			
Agar		15.00			
Final pH (at 25°C) : 5.5 ± 0.2					
<b>Directions :</b>					
Suspend 84.63 grams in 1000 ml distilled water. Heat the medium just to boiling. Sterilize by autoclaving at 15 lbs pressure (121°C) for 10 minutes. AVOID OVERHEATING. Stir the medium while dispensing to prevent settling of calcium carbonate.					
<b>Principle :</b>					
Tomato juice broth which provides nutrients and acid environment for the growth of acidophilic bacteria. Peptonized milk provides lactose as an energy source. The low pH of the medium inhibits bacteria other than acidophilic bacteria. Polysorbate 80 serves as a source of fatty acids. Bromo cresol green acts as a pH indicator. Acid producing bacteria produce a clear yellow halo around the colonies. Other bacteria produce colonies in colours ranging from colourless to yellow green and blue depending on species and strain.					
<b>QC Tests - (I) Dehydrated Medium</b>					
Colour :		Ligth yellow to light green			
Appearance :		Homogeneous Free Flowing powder			
<b>(II) Rehydrated medium</b>					
pH (post autoclaving/heating) :		5.5 ± 0.2			
Colour (post autoclaving/heating) :		Green to light blue			
Clarity (post autoclaving/heating) :		Opaque gel			
<b>(III) Q.C. Test Microbiological</b>					
Cultural characteristics observed after an incubation at 25-30°C for 48-72 hours.					
MICROORGANISM (ATCC )		GROWTH			
Acinetobacter calcoaceticus (23055)		None-poor			
Lactobacillus acidophilus (4356)		luxuriant with clear yellow halo			
Lactobacillus fermentum (9338 )		luxuriant with clear yellow halo			
Lactobacillus leichmannii (4797)		luxuriant with clear yellow halo			
Lactobacillus plantarum(8014)		luxuriant with clear yellow halo			
Proteus vulgaris (13315)		inhibited			
<b>Precautions :</b>		1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials. 3. Cycloheximide is very toxic. Avoid skin contact or aerosol formation and inhalation.			
<b>Limitations :</b>		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.			
<b>Use :</b>		It is used in the brewing industry for the cultivation and identification of brewing bacteria including fastidious type.			
<b>Storage :</b>		Dehydrated medium and prepared medium – Between 2 to 8°C.			
<b>Packing :</b>		500 gm. bottle			
<b>Product profile:</b>	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
<b>B1201</b>	84.63 g/l	5.908 lit	5.5 ± 0.2	Nil	121°C/10 min

**Disclaimer:**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.