

BIOMARK Laboratories-INDIA

www.biomarklabs.com

TECHNICAL SHEET

B128	BUFFERED GLUCOSE BROTH (MR VP MEDIUM)					
Formula						
Ingredients :		gms/lit.				
Buffered peptone		7.00				
Dextrose		5.00				
Dipotassium phosphate		5.00				
Final pH (at 25°C) : 6.9 ± 0.2						
Directions :						
Suspend 17 gms in 1000 ml. distilled water. Heat if necessary to dissolve medium completely. Distribute in 10 ml amounts in tubes. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.						
Principle :						
Buffered peptone and dextrose act as source of carbon, nitrogen, vitamins and minerals. Phosphates buffer the medium.						
QC Tests – (I) Dehydrated Medium						
Colour :		Cream to yellow				
Appearance :		Homogeneous Free Flowing powder				
(II) Rehydrated medium						
pH (post autoclaving/heating) :		6.9 ± 0.2				
Colour (post autoclaving/heating) :		Light yellow				
Clarity (post autoclaving/heating) :		Clear				
(III) Q.C. Test Microbiological						
Cultural characteristics observed after 18 –48 hrs at 30-32°C.						
MICROORGANISM (ATCC)		GROWTH	MR Test	VP Test		
Escherichia coli (25922)		Luxuriant	Positive with bright red color	Negative with no change		
Enterobacter aerogenes (13048)		Luxuriant	Negative reaction (yellow color)	Positive reaction, Eosin pink/red color within 2-5 minutes		
Klebsiella pneumoniae (23357)		Luxuriant	Negative reaction (yellow color)	Positive reaction, Eosin pink/red color within 2-5 minutes		
Precautions :		1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use:		It is recommended for the performance of the Methyl Red and Voges-Proskauer tests in differentiation of the coli-aerogenes group.				
Storage :		Dehydrated medium- below 30°C Prepared medium– Between 2 to 8°C.				
Packing :		500 gm. bottle				
Product profile:		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B128		17 g/l	29.41 L	6.9 ± 0.2	NIL	121°C / 15 minutes

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARK LABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.