

**TECHNICAL SHEET**

<b>B1351</b>	<b>BILE ESCULIN AZIDE BROTH, MODIFIED</b>		
<b>Formula</b>			
<b>Ingredients :</b>	<b>gms/lit.</b>		
Casein enzymic hydrolysate	17.00		
Peptic digest of animal tissue	3.00		
Yeast extract	5.00		
Oxgall	10.00		
Sodium chloride	5.00		
Esculin	1.00		
Ferric ammonium citrate	0.50		
Sodium azide	0.25		
Sodium citrate	1.00		
Final pH (at 25°C) :	7.1 ± 0.2		
<b>Directions :</b>			
Suspend 42.75 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Caution: Sodium azide has a tendency to form explosive metal azides with plumbing materials. It is advisable to use enough water to flush off the disposables.			
<b>Principle :</b>			
Casein enzymic hydrolysate, peptic digest of animal tissue and yeast extract serve as a source of carbon, nitrogen and essential nutrients. Sodium azide inhibits growth of gram-negative organisms and permits the cultivation of Enterococci and group D Streptococci. Oxgall inhibits gram-positive bacteria other than Enterococci. Sodium citrate acts as a buffering agent. Esculin is hydrolysed by Enterococci and group D streptococci to esculin which reacts with ferric ammonium citrate to form dark brown or black complex			
<b>QC Tests - (I) Dehydrated Medium</b>			
Colour :	Cream to yellow		
Appearance :	Homogeneous Free Flowing powder		
<b>(II) Rehydrated medium</b>			
PH (post autoclaving/heating) :	7.1 ± 0.2		
Colour (post autoclaving/heating) :	Amber		
Clarity (post autoclaving/heating) :	Clear to slightly opalescent solution with a bluish tinge.		
<b>(III) Q.C. Test Microbiological</b>			
Cultural characteristics observed after 18 -24 hrs at 35-37°C.			
MICROORGANISM (ATCC )	GROWTH	ESCULIN HYDROLYSIS	
Enterococcus faecalis (29212 )	Luxuriant	+	
Escherichia coli (25922)	None-poor	-	
Staphylococcus aureus (25923 )	Good	-	
Streptococcus pyogenes (19615 )	None - poor	-	
Key : + = blackening of medium - = no change			

Refer Disclaimer overleaf

**TECHNICAL SHEET**

<b>Precautions :</b>	1. For Laboratory Use.				
	2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
	3. HARMFUL. Irritating to eyes, respiratory system and skin. Avoid contact with skin and eyes. Do not breathe dust. Wear suitable protective clothing. Keep container tightly closed. Target organ(s) : Cardiovascular, Lungs, Nerves.				
<b>Limitations :</b>	Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
<b>Use:</b>	It is used to differentiate between Enterococci and Group D Streptococci.				
<b>Storage :</b>	Dehydrated medium-below 30°C Prepared medium- Between 2 to 8°C.				
<b>Packing :</b>	500 gm. bottle				
<b>Product profile:</b>	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
<b>B1351</b>	42.75g/l	11.69 L	7.1 ± 0.2	NIL	121°C / 15 minutes

**Disclaimer:**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.