

BIOMARK Laboratories-INDIA

www.biomarklabs.com

TECHNICAL SHEET

B1375	BUFFERED PEPTONE WATER				
Formula					
Ingredients:		gms/lit.			
Enzymatic digest of casein	10.00				
Sodium chloride		5.00			
Disodium hydrogen phosphate.12H ₂ O		9.00			
Potassium dihydrogen phosphate		1.50			
Final pH (at 25°C) :7.0 ± 0.1					
Directions:					
Suspend 20.07 grams(equivalent weight of dehydrated medium) in 1000 ml distilled water. Heat if necessary to dissolve the medium completely. Dispense as desired and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.					
Principle:					
Buffered Peptone Water containsenzymatic digest of casein as a source of carbon, nitrogen, vitamins and minerals. Sodium Chloride maintains the osmotic balance. Phosphates buffer the medium.					
QC Tests - (I)Dehydrated Medium					
Colour:	Cream to yellow				
Appearance:	Homogeneous Free Flowing powder				
(II)Rehydrated medium					
pH (post autoclaving/heating):	7.0 ± 0.2				
Colour (post autoclaving/heating):	Light yellow to light amber				
Clarity (post autoclaving/heating):	Clear solution				
(III)Q.C. Test Microbiological: ISO 6579 & ISO 21528					
Cultural characteristics observed after 18 -24 hrs at 35-37°C.					
MICROORGANISM (ATCC)	GROWTH				
Salmonella enteritidis (13076)	Luxuriant				
Salmonella typhimurium (14028)	Luxuriant				
Escherichia coli (25922)	Luxuriant				
Escherichia coli (8739)	Luxuriant				
Pseudomonas aeruginosa (27853)	Luxuriant				
Cronobacter sakazakii (29544)	Luxuriant				
Precautions :	1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium. 2. The types and numbers of competing flora in the test sample can affect recovery and may overgrow salmonellae.				
Use :	It is used as pre-enrichment medium for increasing the recovery of injured Salmonella species from foods prior to selective enrichment and isolation.				
Storage :	Dehydrated medium- Below30°C. Prepared mediums- Between 2 - 8°C.				
Packing :	500 gm. bottle				
Product profile:	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B1375	20.07g/l	24.912L	7.0 ± 0.1	NIL	121°C / 15 minutes

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.