#### **BIOMARK Laboratories – INDIA**

# www.biomarklabs.com

## **TECHNICAL SHEET**

B1488 MAC CONKEY BROTI	B1488 MAC CONKEY BROTH PURPLE W/BCP							
Formula								
	jms/lit.							
Peptic digest of animal tissue	20.00							
Lactose	.0.00							
Bile salts	5.00							
Sodium chloride	5.00							
Bromocresol purple (	0.01							
Final pH (at 25°C): 7.4 <u>+</u> 0.2	Final pH (at 25°C): 7.4+ 0.2							
Directions:								
Suspend 40.01 grams in 1000 ml dis	tilled water. Heat if necessary to dissolve the medium							
completely. Dispense into test tubes with	n inverted Durham tubes. Sterilize by autoclaving at 15 lbs							
pressure (121°C) for 15 minutes. Cool the tubes to 45-50°C before inoculation.								
Principle: Peptic digest of animal tissue provides essential growth nutrients. Lactose is the								
fermentable carbohydrate. Bile salts or sodium taurocholate inhibits gram-positive organisms.								
Sodium chloride maintains the osmotic balance of the medium. Bromocresol purple is the pH								
indicator in the medium, which turns yellow under acidic condition. Lactose fermenting organisms								
turn the medium yellow due to the acidity produced on lactose fermentation.								
QC Tests – (I)Dehydrated Medium								
Colour:	Cream to yellow							
Appearance:	Homogeneous Free Flowing powder							
(II)Rehydrated medium								
(11)Renyarated medium								
pH (post autoclaving/heating) :	7.4 ± 0.2							
	7.4 ± 0.2 Purple							

Colour (post autociaving/neating).										
Clarity (post autoclaving/heating):			Clear to slightly opalescent solution in tubes							
(III) Cultural Response			Growth Promotion is carried out in accordance with the							
			harmonized method of IP. For organisms not specified in							
		pharmacopoeia, cultural response was observed after an								
		incubation at 30-35°C for 18-48 hours.								
			Clearly visible growth of microorganism comparable to that previously obtained with previously tested and approved lot							
										of medium occurs at the specified temperature for not more
	than the shortest period of time specifie cfu (at 42-44°C for 24 hours).					1  moculating  <=100				
			No growth of the test microorganism occurs for the specified							
			temperature for not less than longest period of time specified inoculating $>=100$ cfu(at 42-44°C for $>= 48$ hours).							
ultural Decreases		mocu		Loocia(a						
ultural Response										
Cultural characteristics observed										
MICROORGANISM (ATCC)	GROWT	Н	ACID	GAS	INCUBATION	INCUBATION				
					TEMPERATURE	PERIOD				
GROWTH PROMOTING	GROWTH PROMOTING									
Escherichia coli (8739)	luxuriant		+	+	42 -44 °C	<=24 hrs				
Inhibitory										
Staphylococcus aureus (6538) i	inhibited				42 -44 °C	>=48 hrs				
Additional Microbiological test	ting									
	1	~+			30 -35 °C	18 -24 hrs				
Escherichia coli (25922)	Luxuria	nι	+	+	30-35-0	10 -24 115				

Page 01 of 02

### BIOMARK Laboratories – INDIA www.biomarklabs.com TECHNICAL SHEET

	Facharichia a		Luxuriant			30 -35 °C	18 -24 hrs			
	Escherichia d	scherichia coli (NTCC9002)		+	+	30 -35 °C	18 - 24 nrs			
	Enterobacter a	erobacter aerogenes (13048		+	+	30 -35 °C	18 -24 hrs			
	. ,		) Fair to good			30 -35 °C	18 -24 hrs			
			) Inhibited			30 -35 °C	>=48 hrs			
	Key: Acid-+=yellow colour									
Precautions : 1. For Laboratory			y Use.	Use.						
	2. Follow proper, established laboratory procedures in handling and disposing infectious materials.									
Lim	itations :	<b>ations :</b> 1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.								
		<ol> <li>Although MacConkey media are selective primarily for gram – negative enteric bacilli, biochemical and, if indicated, serological testing using pure cultures are recommended for complete identification. Consult appropriate references for further information.</li> </ol>								
Use	:	It is recommended by International Organization for Standardization (ISO), 1990, Draft ISO/ DIS 9308-2 for the presumptive identification of coliforms from water.								
Sto	rage:	Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.								
	king:	500 gm. bottle								
Pro	duct profile:		Quantity on Preparation (500g)	pH (25	°C)	Supplement	Sterilization			
<b>B14</b>	88	40.01 g/L	12.496 L	7.4 <u>+</u> 0.	.2	NIL	121ºC /15 min.			

#### Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

Page 02 of 02