

B151	COOKED MEAT MEDIUM (R.C. MEDIUM)					
Formula						
Ingredients :		gms/lit.				
Meat heart, solids#		98.00				
Proteose peptone		20.00				
Dextrose		2.00				
Sodium chloride		5.00				
#- Equivalent to Beef heart, solids						
Final pH (at 25°C) : 7.2 ± 0.2						
Directions :						
Suspend 12.5 grams in 100 ml distilled water (or suspend 1.25 grams in 10 ml distilled water in test tubes). Mix thoroughly and allow to stand for 15 minutes until all the particles are thoroughly wetted. Dispense into tubes or flasks as desired. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.						
Principle :						
Meat Heart and Proteose Peptone provide the nitrogen, vitamins and amino acids in Cooked Meat Medium. Sodium Chloride maintains the osmotic balance of the medium. The low concentration of Dextrose is sufficient as the energy source, but not high enough to accumulate toxic metabolites. This formulation provides an effective maintenance medium. Solid meat particles provide favorable growth conditions for anaerobes due to the reducing action of -SH (sulfhydryl) groups of muscle protein. Sulfhydryl groups are more accessible in denatured proteins, therefore the use of cooked meat particles is preferred.						
QC Tests - (I) Dehydrated Medium						
Colour :		Brown coloured granules with powder				
Appearance :		Granules with powder				
(II) Rehydrated medium						
pH (post autoclaving/heating) :		7.2 ± 0.2				
Colour (post autoclaving/heating) :		Medium amber				
Clarity (post autoclaving/heating) :		Clear to slightly opalescent supernatant over insoluble granules.				
(III) Q.C. Test Microbiological						
Cultural characteristics observed after 40- 48 hrs. at 35-37°C.						
MICROORGANISM (ATCC)		GROWTH				
Clostridium botulinum (25763)		Luxuriant				
Clostridium perfringens (12924)		Luxuriant				
Clostridium sporogenes (11437)		Luxuriant				
Streptococcus pneumoniae (6303)		Luxuriant				
Enterococcus faecalis (29212)		Luxuriant				
Precautions :		1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :		For cultivation of aerobes and anaerobes particularly pathogenic Clostridia and also for maintenance of stock cultures.				
Storage :		Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
Packing :		500 gm bottle				
Product profile:		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B151	125g/l	4L	7.2 ± 0.2	NIL	121°C / 15 minutes	

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.