

**BIOMARK Laboratories-INDIA**

[www.biomarklabs.com](http://www.biomarklabs.com)

**TECHNICAL SHEET**

<b>B156</b>	<b>CZAPEK DOX AGAR</b>					
<b>Formula</b>						
<b>Ingredients :</b>		<b>gms/lit.</b>				
Sucrose		30.00				
Sodium nitrate		2.00				
Dipotassium phosphate		1.00				
Magnesium sulphate		0.50				
Potassium chloride		0.50				
Ferrous sulphate		0.01				
Agar		15.00				
Final pH (at 25°C) : 7.3 ± 0.2						
<b>Directions :</b>						
Suspend 49.01 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and pour into sterile Petri plates.						
<b>Principle :</b>						
Sucrose is the sole carbon source, and Sodium Nitrate is the sole nitrogen source in Czapek – Dox Agar. Dipotassium Phosphate is the buffering agent, and Potassium Chloride contains essential ions. Magnesium Sulfate and Ferrous Sulfate serves as sources of essential ions. Agar is the solidifying agent in Czapek Dox Agar.						
<b>QC Tests – (I) Dehydrated Medium</b>						
Colour :		Cream to yellow				
Appearance :		Homogeneous Free Flowing powder				
<b>(II) Rehydrated medium</b>						
pH (post autoclaving/heating) :		7.3 ± 0.2				
Colour (post autoclaving/heating) :		Cream to light yellow				
Clarity (post autoclaving/heating) :		Clear to slightly opalescent with a slight precipitate				
<b>(III) Q.C. Test Microbiological</b>						
Cultural characteristics observed after 48-72 hrs. at 25-30°C.						
MICROORGANISM (ATCC )		GROWTH				
Aspergillus niger (16404 )		Luxuriant				
Saccharomyces cerevisiae (9763 )		Luxuriant				
Candida albicans (10231 )		Luxuriant				
<b>Precautions :</b>		1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
<b>Limitations :</b>		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
<b>Use :</b>		For general cultivation of fungi.				
<b>Storage :</b>		Dehydrated medium- below 30°C Prepared medium– Between 2 to 8°C.				
<b>Packing :</b>		500 gm. Bottle				
<b>Product profile:</b>		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
<b>B156</b>		49.01g/l	10.41 L	7.3 ± 0.2	Nil	121°C / 15 minutes

**Disclaimer:**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARK LABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.