

<b>B163</b>	<b>DEOXYCHOLATE CITRATE AGAR</b>					
<b>Formula</b>						
<b>Ingredients :</b>		<b>gms/lit.</b>				
Heart infusion solids		10.00				
Proteose peptone		10.00				
Lactose		10.00				
Sodium citrate		20.00				
Ferric ammonium citrate		2.00				
Sodium deoxycholate		5.00				
Neutral red		0.02				
Agar		13.50				
Final pH (at 25°C) : 7.5 ± 0.2						
<b>Directions :</b>						
Suspend 70.52 grams in 1000 ml of distilled water. Heat to boiling to dissolve the medium completely. DO NOT AUTOCLAVE. Avoid excessive heating as it is detrimental to the medium. Cool to 45-50°C. Mix well and pour into sterile petri plates.						
<b>Principle :</b>						
Heart infusion solids is a source of carbon and nitrogen. Deoxycholate Citrate Agar contains Proteose Peptone as a source of carbon, nitrogen, vitamins and minerals. Lactose is a carbohydrate. Sodium citrate and Sodium deoxycholate inhibit gram positive bacteria, coliforms and Proteus species. Ferric Ammonium Citrate aids in the detection of H <sub>2</sub> S producing bacteria. Neutral Red is a pH indicator. Agar is a solidifying agent.						
In the presence of neutral red, bacteria that ferment lactose produce acid and form red colonies. Bacteria that do not ferment lactose form colorless colonies. If the bacteria produce H <sub>2</sub> S, the colonies will have black centers. The majority of normal intestinal bacteria ferment lactose and do not produce H <sub>2</sub> S (pink-red colonies without black centers). Salmonella and Shigella sp. Do not ferment lactose but Salmonella may produce H <sub>2</sub> S (colourless colonies with or without black centers). Lactose – fermenting colonies may have a zone of precipitation around them caused by the precipitation of deoxycholate in the presence of acid.						
<b>QC Tests - (I)Dehydrated Medium</b>						
	Colour :	Light yellow to pinkish beige				
	Appearance :	Homogeneous Free Flowing powder				
<b>(II)Rehydrated medium</b>						
	pH (post autoclaving/heating) :	7.5 ± 0.2				
	Colour (post autoclaving/heating) :	Reddish orange				
	Clarity (post autoclaving/heating) :	Clear to very slightly opalescent				
<b>(III)Q.C. Test Microbiological</b>						
Cultural characteristics observed after 18-24 hrs.at 35-37°C.						
	MICROORGANISM (ATCC )	GROWTH	COLOUR OF COLONY	H <sub>2</sub> S		
	Salmonella enteritidis (13076 )	Good-luxuriant	Colourless	positive reaction,black centered colonies		
	Salmonella typhimurium (14028)	Good-luxuriant	Colourless	positive reaction,black centered colonies		
	Salmonella Abony (NCTC6017)	Good-luxuriant	Colourless	positive reaction,black centered colonies		
	Shigella flexneri (12022)	Good	Colourless	negative reaction		
	Escherichia coli (25922)	Poor	Pink w/bile ppt.	negative reaction		
	Escherichia coli (8739)	Poor	Pink w/bile ppt.	negative reaction		
	Escherichia coli (NCTC9002)	Poor	Pink w/bile ppt.	negative reaction		
	Streptococcus faecalis (29212 )	Inhibited	-	negative reaction		
	Staphylococcus aureus (25923)	Inhibited	-	negative reaction		
<b>Precautions :</b>		1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
<b>Limitations :</b>		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium. 2. Coliform strains may be encountered that will grow on this medium, making it difficult to detect pathogens. 3. Heavy inocula should be distributed over the entire surface of the medium prevent complete masking of pathogens by coliform organisms.				
<b>Use :</b>		For selective isolation of enteric pathogens especially Salmonella and Shigella spp.				
<b>Storage :</b>		Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
<b>Packing :</b>		500 gm bottle				
<b>Product profile:</b>		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
<b>B163</b>	70.52 g/l	7.09L	7.5 ± 0.2	NIL	DO NOT AUTOCLAVE	

**Disclaimer:**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.