

**BIOMARK Laboratories-INDIA**

[www.biomarklabs.com](http://www.biomarklabs.com)

**TECHNICAL SHEET**

<b>B167</b>	<b>DEXTROSE PEPTONE AGAR</b>					
<b>Formula</b>						
<b>Ingredients :</b>			<b>gms/lit.</b>			
Peptic digest of animal tissue			20.00			
Dextrose			10.00			
Sodium chloride			5.00			
Agar			15.00			
Final pH (at 25°C) : 7.2 ± 0.2						
<b>Directions :</b>						
Suspend 50 gms in 1000 ml. distilled water. Boil to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.						
<b>Principle :</b>						
Peptic digest of animal tissue supplies amino acids, peptides etc. for the growth of the organisms. Dextrose is the readily available energy source for the most of the organisms. The agar medium is also used as an excellent basal agar for the Glucose Blood Agar preparation. In the special petri plates or in deep tubes, it can support good growth of the anaerobic microorganisms.						
<b>QC Tests - (I) Dehydrated Medium</b>						
Colour :			Light yellow			
Appearance :			Homogeneous Free Flowing powder			
<b>(II) Rehydrated medium</b>						
pH (post autoclaving/heating) :			7.2 ± 0.2			
Colour (post autoclaving/heating) :			Light yellow			
Clarity (post autoclaving/heating) :			Clear to slightly opalescent			
<b>(III) Q.C. Test Microbiological</b>						
Cultural characteristics observed after 18 -24 hrs.at 35-37°C.						
MICROORGANISM (ATCC )			GROWTH			
Escherichia coli (25922)			Luxuriant			
Pseudomonas aeruginosa (27853)			Luxuriant			
Staphylococcus aureus (25923)			Luxuriant			
Streptococcus pyogenes (19615)			Luxuriant			
<b>Precautions :</b>		1. For Laboratory Use.				
		2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
<b>Limitations :</b>		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
<b>Use :</b>		For general cultivation of organisms.				
<b>Storage :</b>		Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
<b>Packing :</b>		500 gm bottle				
<b>Product profile:</b>		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
<b>B167</b>		50g/l	10L	7.2 ± 0.2	NIL	121°C / 15 minutes

**Disclaimer:**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications. The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.