## **BIOMARK Laboratories-INDIA** www.biomarklabs.com

## **TECHNICAL SHEET**

FUNGAL BROTH (MYCOLOGICAL BROTH)

B195   FUNGAL BROTH (MYCOLOGICAL BROTH)							
Formula							
		s/lit.					
Pepaic digest of soyabean meal 10.00							
Dextrose		40.0	00				
Final pH (at 25°C): 7.0 <u>+</u> 0.2							
Directions :							
Suspend 50 grams in 1000 ml distilled water. Heat, if necessary, to dissolve the medium completely.							
Dispense as desired. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. For preparing							
selective media, acidify the media upto pH 3.0-4.0 by the addition of two vials of 10% Lactic Acid Solution							
(BF079).							
Principle:							
Papaic digest of soyabean meal in the medium provides nitrogen, vitamins and minerals necessary to							
support bacterial growth. Dextrose is a carbon source required for the growth of fungi.							
QC Tests - (I)Dehydrated Medium							
Colour:			Cream to yellow				
Appearance :			Homogeneous Free Flowing powder				
(II)Rehydrated medium							
pH (post autoclaving/heating) :			$7.0 \pm 0.2$				
Colour (post autoclaving/heating):			Light amber				
Clarity (post autoclaving/heating):			Clear				
(III)Q.C. Test Mi							
Cultural characteristics observed after an incubation at 25-30°C for 48-72 hours(For Trichophyton							
species further incubation may be required for upto 7 days )							
			FUNGAL AGAR WITH LOW pH				
			Luxuriant				
			Luxuriant				
Lactobacillus acidophilus (11506)			Luxuriant				
Staphylococcus aureus (25923)			Inhibited				
Saccharomyces cerevisiae (9763)			Luxuriant				
Saccharomyces uvarum (9080)			Luxuriant				
			Luxuriant				
Precautions:	1. For Laboratory Use.						
	2. Follow proper, established laboratory procedures in handling and disposing of						
	infectious mater						
<b>Limitations :</b> 1. Since the nutritional requirements of organisms vary, some strains may be						ins may be	
	encountered that	countered that fail to grow or grow poorly on this medium.					
Use:	Recommended for cultivation of fungi.						
Storage:	Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.						
Packing: 500 gm. bottle							
	Reconstitution Quantity		on	pH (25°C)	Supplement	Sterilization	
			ion (500g)				
B195	50g/l	10	0.0L	$7.0 \pm 0.2$	NIL	121°C / 15 minutes	
Disclaimer:	1				<u>l</u>		

B195

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

Rev: December 2020