

BIOMARK Laboratories-INDIA

www.biomarklabs.com

TECHNICAL SHEET

B215	KOSER CITRATE MEDIUM					
Formula						
Ingredients :		Gms/lit.				
Sodium ammonium phosphate		1.50				
Monopotassium phosphate		1.00				
Magnesium sulphate		0.20				
Sodium citrate		3.00				
Final pH (at 25°C) : 6.7 ± 0.2						
Directions :						
Suspend 5.7 grams in 1000 ml distilled water. Dispense into tubes and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.						
Principle :						
Koser Citrate Medium is prepared with chemically pure salts and tested to determine that no sources of carbon (other than sodium citrate) or nitrogen (other than ammonium salts) are present. Bacteria that are able to use citrate as their carbon source will grow in the medium and cause turbidity.						
QC Tests – (I) Dehydrated Medium						
Colour :		White to cream				
Appearance :		Homogeneous Free Flowing powder				
(II) Rehydrated medium						
pH (post autoclaving/heating) :		6.7 ± 0.2				
Colour (post autoclaving/heating) :		Colourless				
Clarity (post autoclaving/heating) :		Clear				
(III) Q.C. Test Microbiological						
Cultural characteristics observed after 18- 24 hrs at 35-37°C.						
MICROORGANISM (ATCC)		GROWTH		CITRATE UTILISATION		
Enterobacter aerogenes (13048)		Luxuriant		Positive reaction,turbidity		
Enterobacter cloacae (23355)		Luxuriant		Positive reaction,turbidity		
Escherichia coli (25922)		None- poor		Negative reaction, no turbidity		
Klebsiella pneumoniae (13883)		Luxuriant		Positive reaction,turbidity		
Precautions :		1. For Laboratory Use.				
		2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
		3. IRRITANT. Irritating to eyes, respiratory system and skin. Avoid contact with skin and eyes. Do not breathe dust. Wear suitable protective clothing. Keep container tightly closed.				
Limitations :		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :		For differentiation of Escherichia coli and Enterobacter aerogenes on the basis of citrate utilization.				
Storage :		Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
Packing :		500 gm. bottle				
Product profile:		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B215		5.7g/l	87.71L	6.7 ± 0.2	NIL	121°C / 15 minutes

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.