

TECHNICAL SHEET

B232	LYSINE IRON AGAR				
Formula					
Ingredients :		gms/lit.			
Peptone		5.00			
Yeast extract		3.00			
Dextrose (Glucose)		1.00			
L-Lysine		10.00			
Ferric ammonium citrate		0.50			
Sodium thiosulphate		0.04			
Bromo cresol purple		0.02			
Agar		15.00			
Final pH (at 25°C) : 6.7 ± 0.2					
Directions :					
Suspend 34.56 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Dispense into tubes and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool the tubes in slanted position to form slants with deep butts.					
Principle :					
Lysine Iron Agar contains peptone which provides carbon and nitrogen sources required for good growth of a wide variety of organisms. Yeast Extract provides vitamins and cofactors required for growth, as well as additional sources of nitrogen and carbon. Dextrose is an energy source. L-Lysine Hydrochloride is the substrate used to detect the lysine decarboxylase and lysine deaminase enzymes. Ferric Ammonium Citrate and Sodium Thiosulfate are indicators of hydrogen sulfide production. Bromo Cresol Purple, a pH indicator, is yellow at or below pH 5.2 and purple at or above pH 6.8 Agar is a solidifying agent.					
QC Tests – (I) Dehydrated Medium					
Colour :		Cream to yellow			
Appearance :		Homogeneous Free Flowing powder			
(II) Rehydrated medium					
pH (post autoclaving/heating) :		6.7 ± 0.2			
Colour (post autoclaving/heating) :		Purple			
Clarity (post autoclaving/heating) :		Clear to slightly opalescent			
(III) Q.C. Test Microbiological					
Cultural characteristics observed after 18 - 24 hours at 35 -37°C.					
MICROORGANISM (ATCC)	GROWTH	BUTT	SLANT	H ₂ S	
Citrobacter freundii (8090)	Luxuriant	A	K	+	
Escherichia coli (25922)	Luxuriant	K	K	-	
Proteus mirabilis (25933)	Luxuriant	A	R	+	
Salmonella typhimurium (14028)	Luxuriant	K	K	+	
Shigella flexneri (12022)	Luxuriant	A	K	-	
Salmonella Arizonae (13314)	Luxuriant	K	K	+	
Salmonella Enteritidis (13076)	Luxuriant	K	K	+	
Key : + = blacking of medium					
- = no blacking of medium					
R = deep red. Lysine deamination					
A = acidic, yellow colour					
K = alkaline, purple, no colour change					
Precautions :		1. For Laboratory Use.			
		2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.			

TECHNICAL SHEET

Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
	2. Salmonella paratyphi A, unlike other Salmonella, does not produce lysine decarboxylase and so produces an alkaline slant and an acid butt.				
	3. H ₂ S – producing Proteus species do not blacken the medium. It is, therefore, suggested that Lysine Iron Agar be used in conjunction with Triple Sugar Agar or other media to confirm differentiation.				
	4. The reaction of Morganella morganii may be variable after 24 hours incubation and may require longer incubation.				
Use :	For the differentiation of enteric organisms especially Salmonella Arizonae based on their ability to decarboxylate or deaminate lysine and to form hydrogen sulphide (H ₂ S).				
Storage :	Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
Packing :	500 gm. bottle				
Product profile:	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B232	34.56 g/l	14.46 lit	6.7 ± 0.2	NIL	121°C/15 min