

BIOMARK Laboratories – INDIA

www.biomarklabs.com

TECHNICAL SHEET

B238	Mac Conkey Agar w/0.15% Bile Salts, CV & NaCl
Formula	
Ingredients :	gms/lit.
Peptone	1.50
Tryptone	1.50
Gelatin peptone	17.00
Lactose	10.00
Bile Salts	1.50
Sodium chloride	5.00
Crystal violet	0.001
Neutral red	0.03
Agar	15.00
Final pH (at 25°C) : 7.1± 0.2	
Directions :	
Suspend 51.53 grams in 1000 ml distilled water. Heat to boiling with gentle swirling to dissolve the agar completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Avoid overheating. Cool to 45 - 50°C and pour into sterile Petri plates. The surface of the medium should be dry when inoculated.	
Principle: Peptone, Tryptone and gelatin peptone are sources of nitrogen, carbon, long chain amino acids and other nutrients. Lactose is a fermentable carbohydrate; Sodium chloride maintains the osmotic equilibrium. Bile salts and crystal violet are selective agents that inhibit growth of gram-positive organisms. Neutral red is the pH indicator dye. Lactose-fermenting strains grow as red or pink colonies and may be surrounded by a zone of acid precipitated bile. The red colour is due to production of acid from lactose, absorption of neutral red and a subsequent colour change of the dye when the pH of medium falls below 6.8. Lactose non-fermenting strains, such as Shigella and Salmonella are colourless, transparent and typically do not alter appearance of the medium.	

QC Tests – (I) Dehydrated Medium		
Colour :	Light yellow to pink	
Appearance :	Homogeneous Free Flowing powder	
(II) Rehydrated medium		
pH (post autoclaving/heating) :	7.1 ± 0.2	
Colour (post autoclaving/heating) :	Red with purplish tinge	
Clarity (post autoclaving/heating) :	Clear to slightly opalescent	
(III) Q.C. Test Microbiological		
Cultural response was observed after an incubation at 30-35°C for 18-72 hours.		
MICROORGANISM (ATCC)	GROWTH	COLOUR OF COLONY
Corynebacterium diphtheriae type gravis	Inhibited	--
Enterobacter aerogenes (13048)	Luxuriant	Pink to red
Escherichia coli (25922)	Luxuriant	Pink to red with bile precipitate
Escherichia coli (NCTC 9002)	Luxuriant	Pink to red with bile precipitate
Escherichia coli (8739)	Luxuriant	Pink to red with bile precipitate
Proteus vulgaris (13315)	Luxuriant	Colourless
Salmonella enteritidis (13076)	Luxuriant	Colourless
Salmonella typhi (6539)	Luxuriant	Colourless
Salmonella paratyphi A	Luxuriant	Colourless
Salmonella paratyphi B	Luxuriant	Colourless
Salmonella Typhimurium (14028)	Luxuriant	Colourless
Salmonella Abony (NCTC6017)	Luxuriant	Colourless
Shigella flexneri (12022)	Fair to good	Colourless
Enterococcus faecalis (29212)	Fair to good	Colourless to pink

Refer disclaimer Overleaf

Page 01 of 02

Rev: December 2020

BIOMARK Laboratories – INDIAwww.biomarklabs.com**TECHNICAL SHEET**

	Staphylococcus aureus (25923)	Inhibited	--		
	Staphylococcus aureus (6538)	Inhibited	--		
	Staphylococcus epidermidis (12228)	Inhibited	--		
Precautions :	<ol style="list-style-type: none"> 1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials. 				
Limitations :	<ol style="list-style-type: none"> 1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium. 2. Although MacConkey media are selective primarily for gram – negative enteric bacilli, biochemical and, if indicated, serological testing using pure cultures are recommended for complete identification. Consult appropriate references for further information. 3. Due to the selective properties of MacConkey Agar CV, some strains of gram – negative enteric bacilli may be encountered that fail to grow or grow poorly on this medium. Some strains of gram – positive organisms may be encountered that are not inhibited or only partially inhibited on this medium ; some strains of enterococci may grow on MacConkey Agar CV after prolonged incubation. 4. Incubation of MacConkey Agar plates under increased CO₂ has been reported to reduce the growth and recovery of a number of strains of gram – negative bacilli. 5. For optimal performance, plates prepared from MacConkey Agar CV should be incubated under aerobic conditions. 				
Use :	For selective isolation and differentiation of coliform organisms and other enteric pathogens.				
Storage :	Dehydrated medium- below 30°C Prepared medium– Between 2 to 8°C.				
Packing :	500 gm. bottle				
Product profile:	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B238	51.53 g/L	9.703L	7.1± 0.2	NIL	121°C /15 min.

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.