

TECHNICAL SHEET

B271	SCHAEDLER BROTH	
Formula		
Ingredients :	gms/lit.	
Casein enzymic hydrolysate	5.67	
Proteose peptone	5.00	
Soya peptone	1.00	
Yeast Extract	5.00	
Dextrose	5.83	
Sodium chloride	1.67	
Dipotassium hydrogen phosphate	0.83	
Tris hydroxymethyl aminomethane	3.00	
L-Cystine	0.40	
Hemin	0.01	
Final pH (at 25°C) : 7.6 ± 0.2		
Directions :		
Suspend 28.41 grams in 1000 ml distilled water. If desired 0.02-0.2% Agar can be added. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C and aseptically add 5% sterile defibrinated blood if desired. Mix well and dispense into tubes or flasks as desired. Avoid overheating and photooxidation of the medium as it will retard the growth of bacteria.		
Principle :		
Schaedler broth is highly nutritious medium due to casein enzymic hydrolysate, proteose peptone, soya peptone and yeast extract. Dextrose is a carbon source, and Tris (Hydroxymethyl) amino methane is used to buffer the medium.		
QC Tests - (I) Dehydrated Medium		
Colour :	Cream to yellow	
Appearance :	Homogeneous Free Flowing powder	
(II) Rehydrated medium		
pH (post autoclaving/heating) :	7.6 ± 0.2	
Colour (post autoclaving/heating) :	Light amber	
Clarity (post autoclaving/heating) :	Clear to slightly opalescent	
(III) Q.C. Test Microbiological		
Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours under anaerobic condition.		
MICROORGANISM (ATCC)	GROWTH	
Bacteroides fragilis (25285)	Luxuriant	
Clostridium butyricum (9690)	Luxuriant	
Clostridium perfringens (12924)	Luxuriant	
Clostridium sporogenes (11437)	Luxuriant	
Streptococcus pyogenes (19615)	Luxuriant	
Escherichia coli (25922)	Inhibited	

Refer disclaimer Overleaf

TECHNICAL SHEET

Precautions :	1. For Laboratory Use.				
	2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
	2. Clinical specimens must be obtained properly and transported to the laboratory in a suitable anaerobic transport container.				
	3. The microbiologist must be able to verify quality control of the medium and determine whether the environment is anaerobic.				
	4. The microbiologist must perform aerotolerance testing on each isolate recovered to ensure that the organism is an anaerobe.				
Use:	For cultivation of wide variety of microorganisms particularly from anaerobic blood cultures.				
Storage:	Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
Packing:	500 gm. bottle				
Product profile:	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B271	28.41g/l	17.599L	7.6 ± 0.2	5% sterile defibrinated blood if desired	121°C / 15 minutes

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.