

**TECHNICAL SHEET**

<b>B287</b>	<b>Phenol Red Inositol Broth</b>			
<b>Formula</b>				
<b>Ingredients :</b>		<b>gms/lit.</b>		
Proteose peptone		10.00		
Meat Extract B#		1.00		
Sodium chloride		5.00		
Phenol red		0.018		
Inositol		5.00		
#- Equivalent to Beef extract				
Final pH (at 25°C) : 7.4 ± 0.2				
<b>Directions :</b>				
Suspend 21.02 grams in 1000 ml distilled water, mix well. Heat if necessary, to ensure complete dissolution. Distribute in fermentation tubes (tubes containing inverted Durham's tubes). Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C.				
<b>Principle :</b>				
Proteose peptone and Meat Extract B serve as sources for carbon and nitrogen. Sodium chloride is the osmotic stabilizer. Phenol red is the pH indicator, which turns yellow at acidic pH i.e. on fermentation of inositol. Gas formation is seen in Durham's tubes.				
<b>QC Tests – (I) Dehydrated Medium</b>				
Colour :		Light yellow to Pink		
Appearance :		Homogeneous Free Flowing powder		
<b>(II) Rehydrated medium</b>				
pH (post autoclaving/heating) :		7.4 ± 0.2		
Colour (post autoclaving/heating) :		Red		
Clarity (post autoclaving/heating) :		Clear		
<b>(III) Q.C. Test Microbiological</b>				
Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours (longer if necessary).				
MICROORGANISM (ATCC )	GROWTH	ACID	GAS	
Citrobacter freundii (8090)	Luxuriant	-	-	
Enterobacter aerogenes (13048)	Luxuriant	+	+	
Escherichia coli (25922)	Luxuriant	-	-	
Klebsiella pneumoniae (13883)	Luxuriant	+	+	
Proteus vulgaris (13315)	Luxuriant	-	-	
Salmonella typhimurium (14028)	Luxuriant	+	+	
Salmonella typhi (6539)	Luxuriant	-	-	
Serratia marcescens (8100)	Luxuriant	+	+	
Shigella flexneri (12022)	Luxuriant	-	-	
Key : - = negative reaction, no colour change or red. + = positive reaction, yellow colour				
<b>Precautions :</b>		1. For Laboratory Use.		
		2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.		
<b>Limitations :</b>		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.		

**TECHNICAL SHEET**

	2. The addition of some carbohydrates to the basal medium may cause an acid reaction. To restore the original pH (and colour of the medium), add 0.1 N sodium hydroxide on a drop - by - drop basis. Take care not to make the medium too alkaline, which would prevent fermentation from occurring within the usual incubation period.				
	3. To ensure accuracy of interpretation, uninoculated control tubes and/or inoculated Phenol Red Broth Base control tubes should be run in parallel with the fermentation tests.				
<b>Use :</b>	For Inositol fermentation studies of microorganisms.				
<b>Storage :</b>	Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
<b>Packing :</b>	500 gm. bottle				
<b>Product profile:</b>	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
<b>B287</b>	21.02 g/l	23.786 L	7.4 ± 0.2	Nil	121°C / 15 minutes

**Disclaimer:**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications. The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.