

B317	SELENITE BROTH (SELENITE F BROTH)					
Formula						
Ingredients :			gms/lit.			
Casein enzymichydrolysate			5.00			
Lactose			4.00			
Sodium phosphate			10.00			
Sodium hydrogen selenite			4.00			
Final pH (at 25°C) : 7.0 ± 0.2						
Directions :						
Suspend 23 gms in 1000 ml. distilled water. Mix well. Warm to dissolve the medium completely. Distribute in sterile test tubes. Sterilize in a boiling water bath or free flowing Steam for 10 minutes. DO NOT AUTOCLAVE. Excessive heating is detrimental. Discard the prepared medium if large amount of selenite is reduced. (indicated by red precipitate at the Bottom of tube/bottle)						
Principle :						
Casein enzymic hydrolysate provides the nitrogen, vitamins and amino acids in Selenite Broth. Lactose is a fermentable carbohydrate. Selenite is reduced by organism growth. A rise in pH decrease the selective activity of the selenite. The acid produced by lactose fermentation helps to maintain a neutral pH. Sodium selenite inhibits the growth of gram – positive bacteria and many gram – negative bacteria. Sodium phosphate is a buffering agent.						
QC Tests – (I)Dehydrated Medium						
Colour :		Light yellow				
Appearance :		Homogeneous Free Flowing powder				
(II)Rehydrated medium						
pH (post autoclaving/heating) :		7.0 ± 0.2				
Colour (post autoclaving/heating) :		Very light amber to light yellow				
Clarity (post autoclaving/heating) :		Clear				
(III)Q.C. Test Microbiological						
Cultural characteristics observed when sub cultured on MacConkey Agar(B238) after an incubation at 35-37°C for 18-24 hours.						
MICROORGANISM (ATCC)		RECOVERY		COLOUR OF COLONY*		
Salmonella typhimurium (14028)		Good -Luxuriant		Colourless		
Salmonella cholerasuis (12011)		Good - Luxuriant		Colourless		
Salmonella typhi (6539)		Good - Luxuriant		Colourless		
Escherichia coli (25922)		No increase in numbers		White to pink with bile precipitate		
Key: * = when sub cultured on MacConkey Agar(B238).						
Precautions :		1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials. 3. VERY TOXIC. Fatal if inhaled or swallowed. Irritating to eyes, respiratory system and skin. Avoid contact with skin and eyes. Do not breathe dust. Wear suitable protective clothing. Keep container tightly closed.				
Limitations :		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :		For isolation of Salmonella species from faeces urine or other pathological materials.				
Storage :		Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
Packing :		500 gm. bottle				
Product profile:		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B317	23g/l	21.739L	7.0 ± 0.2	NIL	DO NOT AUTOCLAVE.	