

<b>B427</b>	<b>BILE SALT AGAR</b>					
<b>Formula</b>						
<b>Ingredients:</b>			<b>gms/litre</b>			
Peptic digest of animal tissue			10.00			
Meat extract			5.00			
Sodium chloride			5.00			
Sodium taurocholate			5.00			
Agar			18.00			
Final pH (at 25°C): 8.2 ± 0.2						
<b>Directions:</b>						
Suspend 43 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour into sterile Petri plates						
<b>Principle:</b>						
Peptic digest of animal tissue and meat extract which provide nitrogenous compounds and other essential nutrients for the growth of enteric bacilli. Sodium taurocholate inhibits contaminating gram-positive organisms. Sodium chloride maintains the osmotic balance of the medium.						
<b>QC Tests – (I) Dehydrated Medium</b>						
Colour:		Cream to yellow				
Appearance:		Homogeneous Free Flowing powder				
<b>(II) Rehydrated medium</b>						
pH (post autoclaving/heating) :		8.2 ± 0.2				
Colour (post autoclaving/heating):		Light amber				
Clarity (post autoclaving/heating):		Clear to slightly opalescent				
<b>(III) Q.C. Test Microbiological</b>						
Cultural characteristics observed after 18-24 hours at 35-37°C.						
ORGANISM (ATCC)		GROWTH				
Enterobacter aerogenes (13048)		luxuriant				
Escherichia coli (25922)		luxuriant				
Staphylococcus aureus (25923)		inhibited				
Salmonella Typhi (6539)		luxuriant				
Vibrio cholerae (15748)		luxuriant				
<b>Precautions :</b>		1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
<b>Limitations :</b>		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
<b>Use:</b>		It is used for isolation and enumeration of bile tolerant enteric bacilli.				
<b>Storage:</b>		Dehydrated medium- below 30°C Prepared medium– Between 2 to 8°C.				
<b>Packing:</b>		500 gm. bottle				
<b>Product profile:</b>		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
<b>B427</b>	43 g/l	11.62L	8.2± 0.2	None	121°C / 15 minutes	

**Disclaimer:**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARK LABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.