

B517	ORANGE SERUM BROTH				
Formula					
Ingredients :					
		gms/lit.			
Casein enzymic hydrolysate		10.00			
Yeast extract		3.00			
Dextrose		4.00			
Dipotassium phosphate		2.50			
Orange serum(solids from 200ml)		9.00			
Final pH (at 25°C) : 5.5 ± 0.2					
Directions :					
Suspend 28.5 grams in 1000 ml distilled water. Heat if necessary to dissolve the medium completely. Dispense as desired. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. AVOID OVERHEATING.					
Principle :					
Casein enzymic hydrolysate provides essential nitrogenous nutrients while dextrose serves as the fermentable carbohydrate and energy source. Yeast extract supplies B- complex vitamins, which stimulate growth. Orange serum provides an optimal environment for the recovery of acid tolerant microorganisms from citrus fruit products.					
QC Tests – (I)Dehydrated Medium					
	Colour :	Cream to yellow			
	Appearance :	Homogeneous Free Flowing powder			
(II)Rehydrated medium					
	pH (post autoclaving/heating) :	5.5 ± 0.2			
	Colour (post autoclaving/heating) :	Medium to dark amber			
	Clarity (post autoclaving/heating) :	Clear to slightly opalescent			
(III)Q.C. Test Microbiological					
Cultural characteristics observed after 40 – 48 hrs at 35- 37°C.					
	MICROORGANISM (ATCC)	GROWTH			
	Aspergillus niger (16404)	Good to luxuriant			
	Lactobacillus fermentum (9338)	Good to luxuriant			
	Lactobacillus acidophilus (4356)	Good to luxuriant			
	Leuconostoc mesenteroides (12291)	Good to luxuriant			
	Saccharomyces cerevisiae (9763)	Good to luxuriant			
	Candida albicans (10231)	Good to luxuriant			
Precautions :	1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :	For cultivation and enumeration of microorganisms associated with spoilage of citrus products, cultivation of Lactobacilli and other aciduric organisms and pathogenic fungi.				
Storage :	Dehydrated medium and prepared medium– Between 2 to 8°C.				
Packing :	500 gm. bottle				
Product profile:	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B517	28.5 g/l	17.54L	5.5 ± 0.2	Nil	121°C/15 min.

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.