

TECHNICAL SHEET

B534	STAPHYLOCOCCUS AGAR NO.110			
Formula				
Ingredients :		gms/lit.		
Casein enzymichydrolysate		10.00		
Yeast extract		2.50		
Gelatin		30.00		
Lactose		2.00		
D-Mannitol		10.00		
Sodium chloride		75.00		
Dipotassium phosphate		5.00		
Agar		15.00		
Final pH (at 25°C) : 7.0± 0.2				
Directions :				
Suspend 149.5 gms in 1000ml. distilled water. Mix thoroughly. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Resuspend the precipitate by gentle agitation to avoid bubbles and pour the plates while the medium is hot. Alternatively, cool the medium to 45-50°C and add blood or egg yolk if desired. Staphylococcus Agar No.110 may also be used without sterilization; it should be boiled for 5 minutes and used at once.				
Principle :				
Staphylococcus Agar No.110 is formulated as described by Chapman (1,2,3) for selective isolation and enumeration of Staphylococci from clinical as well as nonclinical specimens. These media are recommended APHA (5). The addition of blood in the medium enables to study haemolytic reaction (6) and with egg yolk enables to study lecithinase production by Staphylococcus aureus (7). These media are selective due to high salt concentration and differential on the basis of ability of organisms to ferment the mannitol, pigment production and gelatin liquefaction. These media are very nutritive as they contain casein enzymichydrolysate, yeast extract which provide essential growth factors like vitamins, nitrogen, carbon compounds, sulphur and trace nutrients etc. to the organisms. High concentration of sodium chloride inhibits many bacterial species except staphylococci. Sodiumazide inhibits gram-negative organisms. Mannitol fermentation can be visualized as yellow colouration by addition of a few drops of bromothymol blue to the areas of the plates from where colonies have been removed. Gelatin liquefaction can be seen when the plates are flooded with a saturated aqueous solution of ammonium sulphate. Enterococcus faecalis may grow on these media as small colonies with little mannitol fermentation (8).				
QC Tests – (I) Dehydrated Medium				
Colour :		Light Yellow		
Appearance :		Homogeneous Free Flowing powder		
(II) Rehydrated medium				
pH (post autoclaving/heating) :		7.0 ± 0.2		
Colour (post autoclaving/heating) :		Light amber		
Clarity (post autoclaving/heating) :		Opalescent		
(III) Q.C. Test Microbiological				
Cultural characteristics observed after 48 hrs at 35 - 37°C.				
MICROORGANISM (ATCC)	GROWTH	PIGMENT PRODUCTION	GELATINASE PRODUCTION	MANNITOL FERMENTATION
Staphylococcus aureus (25923)	Luxuriant	+	+	+
Staphylococcus epidermidis (12228)	Luxuriant	-	+	V
Enterococcus faecalis (29212)	None-poor	-	V	±
Escherichia coli (25922)	Inhibited	-	-	-
Key : + = positive reaction - = negative reaction V = variable reaction ± = slight reaction				

BIOMARK Laboratories-INDIAwww.biomarklabs.com**TECHNICAL SHEET**

Precautions :	1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :	B534: For selective isolation and differentiation of pathogenic Staphylococci.				
Storage :	Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
Packing :	500 gm bottle				
Product profile:	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B534	149.5g/l	3.344L	7.0 ± 0.2	blood or egg yolk	121 ⁰ C / 15 minutes

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.