

**TECHNICAL SHEET**

<b>B540</b>	<b>GLUCOSE BROTH</b>					
<b>Formula</b>						
<b>Ingredients :</b>		<b>gms/lit.</b>				
Casein enzymic hydrolysate		10.00				
Glucose		5.00				
Sodium chloride		5.00				
Final pH (at 25°C) : 7.3 ± 0.2						
<b>Directions :</b>						
Suspend 20 grams in 1000 ml distilled water. Heat if necessary, to dissolve the medium completely. Dispense in tubes containing inverted Durhams tubes. Sterilize by autoclaving at 118°C for 15 minutes.						
<b>Principle :</b>						
Casein enzymic hydrolysate and glucose serve as sources of essential nutrients and energy respectively to support the growth of many fastidious organisms. The casein enzymic hydrolysate used is free of carbohydrates and glucose acts as source of energy by being the only fermentable carbohydrate. The broth gives rapid growth and hastens the early development of injured cells. Sodium chloride maintains the osmotic equilibrium.						
<b>QC Tests - (I) Dehydrated Medium</b>						
Colour :		Light yellow				
Appearance :		Homogeneous Free Flowing powder				
<b>(II) Rehydrated medium</b>						
pH (post autoclaving/heating) :		7.3 ± 0.2				
Colour (post autoclaving/heating) :		Light yellow				
Clarity (post autoclaving/heating) :		Clear				
<b>(III) Q.C. Test Microbiological</b>						
Cultural characteristics observed after 18 –24 hrs. at 35-37°C.						
MICROORGANISM (ATCC )		GROWTH		GAS		
Escherichia coli (25922)		Luxuriant		+		
Salmonella typhi ( 6539 )		Luxuriant		-		
Shigella flexneri (12022)		Luxuriant		-		
Staphylococcus aureus(25923)		Luxuriant		-		
Staphylococcus epidermidis (12228)		Luxuriant		-		
Streptococcus pyogenes (19615)		Luxuriant		-		
<b>Precautions :</b>		1. For Laboratory Use.				
		2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
<b>Limitations :</b>		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
<b>Use :</b>		For study of dextrose fermentation whereas pH indicator is not desired.				
<b>Storage :</b>		Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
<b>Packing :</b>		500 gm bottle				
<b>Product profile:</b>		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
<b>B540</b>		20g/l	25 L	7.3 ± 0.2	nil	118°C / 15 minutes

**Disclaimer:**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.