BIOMARK Laboratories-INDIA www.biomarklabs.com TECHNICAL SHEET

B600						EROFERM SCREE	
Formula			DILACIUS	ACIL			
Ingredients :		gms/lit.					
Dextrose	20.00						
Proteose peptone	10.00						
Yeast extract	5.00						
Sodium acetate	5.00						
2-Phenylethyl alc	3.00						
Ammonium citrat	2.00						
Dipostassium pho	2.00						
Magnessium sulpl	0.10						
Manganese sulph	0.10						
Bromo cresol gree	0.03						
Cycloheximide	0.004						
		2					
Final pH (at 25°C) : 4.3 <u>+</u> 0.2							
Directions :							
Suspend 47.2 grams in 1000 ml distilled water containing 1 ml polysorbate 80. Mix thoroughly and							
dispense in tubes containing inverted Durhams tubes. Sterilize by autoclaving at 15 lbs pressure (121°C)							
for 15 minutes. If necessary, adjust the pH with glacial acetic acid after sterilization.							
Warning: Cycloheximide is very toxic. Avoid skin contact or aerosol formation and inhalation.							
Principle :							
Proteose peptone and dextrose supply nitrogen, carbon and other elements essential for the growth of							
Lactobacilli. Polysorbate 80 a mixture of oleic esters, supplies fatty acids required by Lactobacilli.							
Ammonium citrate, Sodium acetate, 2-phenylethyl alcohol and Cycloheximide inhibit gram – negative							
organisms, molds and certain gram - positive bacteria. Certain yeasts are suppressed because of							
presence of Cycloheximide.							
QC Tests - (I)D	ehydrated Med	ium					
Colour :		Light yellow to bluish grey					
Appearance :		Homogeneous Free Flowing powder					
(II)Rehydrated							
pH (post auto	:	4.3 ± 0.2					
Colour (post a	ng):	Green					
Clarity (post a		Clear to slightly opalescent					
(III) Q.C. Test N	5/						
Cultural characteristics observed after upto 3 days at 35-37°C with 5 – 10% CO ₂ .							
MICROORGAN			GROWT			2070 002	
Lactobacillus p)	Luxuria					
Lactobacillus f	-	Luxuria					
Lactobacillus acidophilus (4356)			Luxuria				
		5)					
Lactobacillus casei (9595) Luxuriant							
Precautions :	1. For Laboratory Use.						
	2. Follow proper, established laboratory procedures in handling and disposing of						
infectious materials. Limitations : 1. Since the nutritional requirements of organisms vary, some strains may be							
							is may be
	encountered that fail to grow or grow poorly on this medium.						
Use :	For the isolation and cultivation of Lactobacillus species from salad dressings.						
Storage :	Dehydrated medium and prepared medium – Between 2 to 8°C. 500 gm. bottle						
Packing :	T-						
Product profile:	Reconstitution	Quantity		рН	(25°C)	Supplement	Sterilization
			ion (500g)				
B600	47.2g/l	10	.593L	4.3	3 ± 0.2	Polysorbate 80	121°C / 15 minutes
 Consultantia transmission Operational 							

Refer disclaimer Overleaf

Page 01 of 02

Disclaimer:

Rev: December 2020

BIOMARK Laboratories-INDIA www.biomarklabs.com TECHNICAL SHEET

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

Page 02 of 02