

BIOMARK Laboratories-INDIA

www.biomarklabs.com

TECHNICAL SHEET

B638	MOTILITY NITRATE MEDIUM, BUFFERED					
Formula						
Ingredients :		gms/lit.				
Peptic digest of animal tissue		5.00				
Meat Extract B#		3.00				
Galactose		5.00				
Potassium nitrate		1.00				
Disodium phosphate		2.50				
Agar		3.00				
#- Equivalent to Beef extract						
Final pH (at 25°C) :		7.3 ± 0.2				
Directions :						
Suspend 19.5 gms. in 1000 ml distilled water containing 5ml. glycerol. Heat to boiling to dissolve the medium completely. Dispense in test tubes to make them half full. Sterilize by autoclaving at 15 lbs pressure (121° C) for 15 minutes. Cool quickly in cool running water and allow the tubed medium to solidify in an upright position.						
Principle :						
Peptic digest of animal tissue and Meat Extract B galactose provide essential nutrients for growth. Potassium nitrate is the substrate for nitrate reduction which is detected with the help of two reagents, viz. sulfanilic acid (1 gm in 125 ml 5 N acetic acid) and N - (1-naphthyl) ethylene diamine dihydrochloride (0.25 gm in 200 ml 5N acetic acid). The presence of less quantity of agar in the medium makes it semisolid which allows detection of motility.						
QC Tests – (I) Dehydrated Medium						
Colour :		Light to medium yellow				
Appearance :		Homogeneous Free Flowing powder				
(II) Rehydrated medium						
pH (post autoclaving/heating) :		7.3 ± 0.2				
Colour (post autoclaving/heating) :		Light amber				
Clarity (post autoclaving/heating) :		Clear to slightly opalescent gel forms in tubes as butt				
(III) Q.C. Test Microbiological						
Cultural characteristics observed after 24- 48 hours at 35-37°C.						
MICROORGANISM (ATCC)		MOTILITY		NITRATE REDUCTION		
Clostridium perfringens (12924)		-		+		
Clostridium absonum (27555)		w		±		
Key : + = red-violet colour, ± weak or absent						
- = growth along stabline, w = weakly motile						
Precautions :		1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :		For isolation and detection of Clostridium perfringens on the basis of motility and nitrate test.				
Storage :		Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
Packing :		500 gm. bottle				
Product profile:		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B638		19.5 g/l	25.64L	7.3 ± 0.2	glycerol	121°C / 15 minutes

Refer disclaimer Overleaf

Rev: December 2020

BIOMARK Laboratories-INDIA
www.biomarklabs.com
TECHNICAL SHEET

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.