

TECHNICAL SHEET

B717	SELENITE BROTH BASE W/O SELENITE					
Formula						
Ingredients :			gms/lit.			
Casein enzymic hydrolysate			5.00			
Lactose			4.00			
Sodium phosphate			10.00			
Final pH (at 25°C) : 7.0 ± 0.2						
Directions :						
Suspend 4 gram of sodium hydrogen selenite in 1000 ml. distilled water. Add 19 grams of B717. Mix well. Warm to dissolve the medium completely. Distribute in sterile test tubes. Sterilize in a boiling water bath or free flowing Steam for 10 minutes. DO NOT AUTOCLAVE. Excessive heating is detrimental. Discard the prepared medium if large amount of selenite is reduced. (indicated by red precipitate at the Bottom of tube/bottle)						
Principle :						
Casein enzymic hydrolysate provides the nitrogen, vitamins and amino acids in Selenite Broth. Lactose is a fermentable carbohydrate. Selenite is reduced by organism growth. A rise in pH decrease the selective activity of the selenite. The acid produced by lactose fermentation helps to maintain a neutral pH. Sodium selenite inhibits the growth of gram – positive bacteria and many gram – negative bacteria. Sodium phosphate is a buffering agent.						
QC Tests – (I) Dehydrated Medium						
Colour :			Cream to light yellow			
Appearance :			Homogeneous Free Flowing powder			
(II) Rehydrated medium						
pH (post autoclaving/heating) :			7.0 ± 0.2			
Colour (post autoclaving/heating) :			Cream to yellow			
Clarity (post autoclaving/heating) :			Clear solution without any precipitate			
(III) Q.C. Test Microbiological						
Cultural characteristics observed with added sodium hydrogen Selenite when subcultured on MacConkey Agar (B238) after an incubation at 35-37°C for 18-24 hours.						
MICROORGANISM (ATCC)		RECOVERY		COLOUR OF COLONY*		
Salmonella typhimurium (14028)		Good -Luxuriant		Colourless		
Salmonella cholerasuis (12011)		Good - Luxuriant		Colourless		
Salmonella typhi (6539)		Good - Luxuriant		Colourless		
Escherichia coli (25922)		No increase in numbers		White to pink with bile precipitate		
Key: * = when subcultured on MacConkey Agar(B238)						
Precautions :		1. For Laboratory Use.				
		2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
		3. VERY TOXIC. Fatal if inhaled or swallowed. Irritating to eyes, respiratory system and skin. Avoid contact with skin and eyes. Do not breathe dust. Wear suitable protective clothing. Keep container tightly closed.				
Limitations :		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :		With addition of selenite, it is used for enrichment of Salmonellae from food, dairy products and pathological materials.				
Storage :		Dehydrated medium- below 30°C Prepared medium– Between 2 to 8°C.				
Packing :		500 gm. bottle				
Product profile:		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B717		19+4 g/l	26.315L	7.0 ± 0.2	Sodium hydrogen selenite	DO NOT AUTOCLAVE.

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.