

B763	Mac Conkey Broth (Double Strength)					
Formula						
Ingredients :		gms/lit.				
Peptic digest of animal tissue		40.00				
Lactose		20.00				
Bile Salts		10.00				
Sodium Chloride		10.00				
Neutral red		0.15				
Final pH (at 25°C): 7.4 ± 0.2						
Directions :						
Suspend 80.15 grams in 1000 ml distilled water. Heat if necessary, to dissolve the medium completely. Distribute into test tubes with inverted Durham tubes and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool the tubes before inoculation						
Principle: Peptic digest of animal tissue provides necessary nitrogen source. Lactose serves as the fermentable carbohydrate source. Sodium chloride maintains the osmotic balance of the cells. The selective action of these media is attributed to the presence of bile salts, which are inhibitory to most species of gram-positive bacteria. Gram-negative bacteria usually grow well on these media and are differentiated by their ability to ferment lactose. The medium turns pink in case of lactose fermenters and yellow in case of lactose- nonfermenters, due to neutral red.						
QC Tests – (I) Dehydrated Medium						
Colour :		Light yellow to pink				
Appearance :		Homogeneous Free Flowing powder				
(II) Rehydrated medium						
pH (post autoclaving/heating) :		7.4 ± 0.2				
Colour (post autoclaving/heating) :		Red				
Clarity (post autoclaving/heating) :		Clear				
(III) Q.C. Test Microbiological						
Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.						
MICROORGANISM (ATCC)		GROWTH	ACID	GAS		
Enterobacter aerogenes (13048)		Luxuriant	+	+		
Escherichia coli (25922)		Luxuriant	+	+		
Salmonella cholerasuis (12011)		Fair to good	--	--		
Staphylococcus aureus (25923)		Inhibited	--	--		
Precautions :		1. For Laboratory Use.				
		2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				

Refer disclaimer Overleaf

BIOMARK Laboratories – INDIAwww.biomarklabs.com**TECHNICAL SHEET**

Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium. 2. Although MacConkey media are selective primarily for gram – negative enteric bacilli, biochemical and, if indicated, serological testing using pure cultures are recommended for complete identification. Consult appropriate references for further information.				
Use :	For primary isolation and coliforms from large samples such as water or waste water.				
Storage :	Dehydrated medium- below 30°C Prepared medium– Between 2 to 8°C.				
Packing :	500 gm. bottle				
Product profile:	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B763	80.15 g/L	6.238L	7.4± 0.2	NIL	121°C /15 min.

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications. The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.