

BIOMARK Laboratories-INDIA

www.biomarklabs.com

TECHNICAL SHEET

B774		THIOBACILLUS AGAR				
Formula						
Ingredients :		gms/lit.				
Ammonium sulphate		0.40				
Monopotassium phosphate		4.00				
Calcium chloride		0.25				
Ferrous sulphate		0.01				
Magnesium sulphate		0.50				
Sodium thiosulphate		5.00				
Agar		12.50				
Final pH (at 25°C) : Self						
Directions :						
Suspend 22.7 gms in 1000 ml distilled water. Boil to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes & dispense as desired.						
Principle :						
Samples are inoculated into Thiobacillus Broth, After incubation at 25-30°C for about 7 days or more, turbidity or sulphur precipitation on the surface of the liquid or against the walls of the flasks, indicates growth of bacteria. Isolation is subsequently done on Thiobacillus Agar. Thiobacillus forms small sulphur impregnated colonies with clear zones, indicating & formation from thiosulphate oxidation.						
QC Tests – (I) Dehydrated Medium						
Colour :		Off white				
Appearance :		Homogeneous Free Flowing powder				
(II) Rehydrated medium						
pH (post autoclaving/heating) :		Self				
Colour (post autoclaving/heating) :		Light amber				
Clarity (post autoclaving/heating) :		Clear to slightly opalescent				
(III) Q.C. Test Microbiological						
Cultural characteristics observed after 7 days at 25 - 30°C.						
MICROORGANISM (ATCC)		GROWTH				
Thiobacillus thiooxidans		Good – luxuriant				
Thiobacillus thioparus		Good – luxuriant				
Precautions :		1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :		B774: For isolation and cultivation of Thiobacillus species.				
Storage :		Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
Packing :		500 gm. bottle				
Product profile:		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B774		22.7 g/l	22.02L	Self	NIL	121°C /15 min.