

<b>B966</b>	<b>CAMPYLOBACTER ENRICHMENT BROTH BASE (PRESTON ENRICHMENT BROTH BASE)</b>					
<b>Formula</b>						
<b>Ingredients :</b>		<b>gms/lit.</b>				
Peptic digest of animal tissue		10.00				
Beef extract		10.00				
Sodium chloride		5.00				
Final pH (at 25°C) : 7.5 ± 0.2						
<b>Directions :</b>						
Suspend 12.5 grams in 470 ml distilled water. Heat if necessary, to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to room temperature and aseptically add sterile 25 ml lysed horse blood and reconstituted contents of 1 vial of Campylobacter Supplement IV (Preston Selective Supplement) (BF033). Mix well and dispense as desired.						
<b>Principle :</b>						
Peptic digest of animal tissue and beef extract in the medium provide nitrogen, vitamins and minerals necessary to support bacterial growth. Sodium chloride provides essential ions.						
<b>QC Tests - (I) Dehydrated Medium</b>						
Colour :		Cream to light yellow				
Appearance :		Homogeneous Free Flowing powder				
<b>(II) Rehydrated medium</b>						
pH (post autoclaving/heating) :		7.5 ± 0.2				
Colour (post autoclaving/heating) :		A) Basal medium : light amber B) (After addition of blood yields): Cherry red				
Clarity (post autoclaving/heating) :		A) Clear B) Opaque				
<b>(III) Q.C. Test Microbiological</b>						
Cultural characteristics observed with added 25ml sterile lysed horse blood and Campylobacter Supplement IV (Preston Selective Supplement), (BF033), after an incubation at 42°C for 48 hours (5% O <sub>2</sub> + 10% CO <sub>2</sub> + 85% N <sub>2</sub> ).						
MICROORGANISM (ATCC )		GROWTH				
Campylobacter jejuni (29428)		Good-luxuriant				
Campylobacter coli (33559)		Good-luxuriant				
Campylobacter lari (35221)		Good-luxuriant				
Escherichia coli (25922)		Inhibited				
Proteus mirabilis (25933)		Inhibited				
Staphylococcus aureus (25923)		Inhibited				
Bacillus cereus (10876)		Inhibited				
<b>Precautions :</b>		1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
<b>Limitations :</b>		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
<b>Use :</b>		For selective enrichment and cultivation of Campylobacter species.				
<b>Storage :</b>		Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
<b>Packing :</b>		500 gm bottle				
<b>Product profile:</b>		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
<b>B966</b>		25g/l	20L	7.5 ± 0.2	sterile 25 ml lysed horse blood and reconstituted contents of 1 vial of Campylobacter Supplement IV (Preston)(BF033)	121°C / 15 minutes