

**BA025**

**PICRIC ACID (SATURATED, AQUEOUS)**

**FORMULA**

**Reagents:**

Picric acid	1.18 g
Distilled water	100.00 ml

**Directions:**

1. Take a loopful of fungi culture on clear, dry glass slide.
2. Mix it with a loopful of picric acid.
3. Tease the fungal culture into thin preparation carefully with the help of wire loop.
4. Place coverslip. Wait for 5 minutes.
5. Observe under high power.
- 6.

**Precautions :**

1. For Laboratory Use.
2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.

**Use:** Fungal staining

**Quality Control:**

**Appearance:** Yellow coloured solution.

**Microscopy:**

Fungal spores and hyphae are stained pale to dark yellow.

**Storage:** Below 30°C.

**Packing :** 100 ml

**Disclaimer:**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.