

**BA036**

**Neisser's methylene blue**

**FORMULA**

**Reagent:**

Methylene blue	1.00 g
Glacial acetic acid	50.00 ml
Ethyl alcohol,95%	50.00 ml
Distilled water	100.00 ml

**Directions:**

1. Take a loopful of culture on clear, dry glass slide. Prepare a thin smear.
2. Air dry and fix with gentle heat.
3. Flood slide with Neisser's methylene blue for 3 minutes.
4. Wash off with 1:10 diluted in distilled water Gram's iodine solution and leave some of this solution on the slide for 1 minute.
5. Wash with water.
6. Counter stain with Neutral Red Solution for 3 minutes.
7. Wash with water, air dry.
8. Observe under oil immersion lens.

**Precautions :**

1. For Laboratory Use.
2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.

**Use:** Metachromatic granule staining.

**Quality Control:**

**Appearance:**Blue coloured solution

**Microscopy:**

Metachromatic granules: Deep blue  
Organisms: Pink

**Storage:** Below 30°C.

**Packing:** 100 ml

**Disclaimer:**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.