

TECHNICAL SHEET

BA041

PAPANICOLAU-OG-6

Formula

Reagent:

Orange G	0.30 g
Phosphotungstic acid	0.015 g
Denatured Alcohol	100.00 ml

Directions:

Fixation: Do not allow smears to dry and fix immediately in 95% alcohol for 5-15 min
The smears may be left in the fixative for 3 days, if necessary, but prolonged fixation affects the Staining reaction.

1. Rinse in 70% alcohol, 50% alcohol and distilled water.
2. Stain in Harris hematoxylin (without acetic acid) for 5 -10 minutes.
3. Rinse in distilled water.
4. Rinse 3 or 4 times in 0.5% aqueous solution of hydrochloric acid.
5. Rinse thoroughly in water.
6. Leave for 1 minute in a weak solution of lithium, carbonate (3drops saturated aqueous solution / 100 ml water). Rinse thoroughly in water.
7. Rinse in distilled water, 50% alcohol, 70% alcohol, 80% alcohol and 95% alcohol.
8. Stain for 1 minute in the Papanicolaou Orange G-6 (BA041) solution.
9. Rinse 5-10 times in each of two jars containing 95% alcohol.
10. Stain in Papanicolaou EA36 (BA042) for 2 minutes.
11. Rinse 5-10 times in each of three jars containing,95% alcohol (not the same alcohol that was used after orange G-6 solution).
12. Rinse in absolute alcohol, then in a mixture of equal parts of absolute alcohol and xylene and then in xylene.
13. Mount in any satisfactory neutral medium and observe under microscope

Quality Control:

Appearance: Orange coloured solution clear without any particles.

Microscopy: Nuclei: Blue Cytoplasm: Grey or pale blue

Storage: Below 30°C.

Packing: 100 ml

Warning & Precautions :

In Vitro diagnostic Use only. For professional use only. For Laboratory Use. Read the label before opening the container. Wear protective gloves/protective clothing/ eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets

Use: Staining of malignant cells

TECHNICAL SHEET

BA041

PAPANICOLAU-OG-6

Disposal:

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques.

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.