

TECHNICAL SHEET

BA066 FOLIN AND WU'S PHOSPHATE MOLYBDATE SOLUTION

Formula

Reagent:

Molybdic acid	35.00 g
Sodium tungstate	5.00 g
10% NaOH solution	200.00 ml
Distilled water	500.00 ml
Phosphoric acid, concentrated	125.00 ml

Appearance: Colourless solution

Directions:

Deproteinisation:

1. Take 1.0 ml of blood in a test tube. Add 7.0 ml of distilled water and 1.0 ml of 0.6 N H₂SO₄. Mix well. Keep it for 5-10 minutes. Add 1 ml of 10% Sodium tungstate to precipitate proteins. Mix well and keep for 10 minutes. Centrifuge it or filter.

Reduction and colour development:

1. Take 2.0 ml of the above filtrate in a tube. Add 2ml of reagent (BA065) in a tube. Mix and keep in boiling water bath for 8 minutes. Cool under running tap water. Add 2 ml of reagent (BA066) and dilute with distilled water to 25 ml. Mix well and keep for 5 minutes. Read the colour at 620 nm.

Precautions :

1. For Laboratory Use.
2. Follow proper, established laboratory procedures in handling and disposing infectious materials.

Use: Estimation of blood sugar.

Storage: Below 30°C

Packing: 500 ml

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.