

BIOMARK Laboratories-INDIAwww.biomarklabs.com**TECHNICAL SHEET****BDR 008****Gram Positive ring with 12 antimicrobial discs****SPECIFICATIONS:****COLOUR:** White to off white ring having 12 discs of 6 mm diameter each with abbreviation.**DIRECTIONS:**

1. Prepare sterile plates of Mueller Hinton Agar (B263).
2. Prepare inoculum as per Bauer-Kirby Method-make a pure inoculum of culture. Inoculate pure single colony of test organism – E. coli (ATCC 25922) .S.aureus (25923) and ps.aeruginosa (27953), in 5ml Pre tested sterile SCDM (B040) in a test tube & incubate at 36 to 37 °C for 2 hours. Compare the inoculum turbidity with that of standard 0.5McFARLAND STANDARD (BA 132).
3. Dip a sterile swab into the standardized inoculum and press against the wall of the test tube to remove excess liquid.
4. Swab the plates by rotating in all directions till the plate becomes dry.
5. Place the antimicrobial ring on the swabbed plates aseptically with the help of alcohol dipped flame sterilized forcep. Press the ring properly on the plates.
6. Incubate the plates at 35 C - 37°C for 18- 24 hours.
7. Measure the zones showing complete inhibition and record the diameters of the zones to the nearest millimeter using a calibrated instrument like zone scales.(BL/AS/01)

RESULTS:

Average diameter of zone of inhibition is observed on Mueller Hinton Agar (B263) after 18-24 hours incubation at 37°C for standard cultures.

Name BDR008 hospitest GRAM POSITIVE 06.09.22	Resistant Less than or =	Intermediate	Sensitive More than or =
Ampicillin (10mcg) Amp EC (ATCC 25922) SA (ATCC 25923) PS (ATCC 27853)	13 28 --	14-16 -- --	17 29 --
Meropenem (10mcg) mem EC (ATCC 25922) SA (ATCC 25923) PS (ATCC 27853)	19 13 15	20-22 14-15 16-18	23 16 19
Erythromycin (5mcg) ery EC (ATCC 25922) SA (ATCC 25923) PS (ATCC 27853)	-- -- --	-- -- --	-- -- --
Tetracyclin (30) tet EC (ATCC 25922) SA (ATCC 25923) PS (ATCC 27853)	11 14 --	12-14 15-18 --	15 19 --
Co trimaxoxazole 25mcg) cot EC (ATCC 25922) SA (ATCC 25923) PS (ATCC 27853)	10 10 --	11-15 11-15 --	16 16 --
Cefuroxime (10mcg) crx EC (ATCC 25922) SA (ATCC 25923) PS (ATCC 27853)	-- -- --	-- -- --	-- -- --
Gentamycin (10mcg) gen EC (ATCC 25922) SA (ATCC 25923) PS (ATCC 27853)	12 12 12	13-14 13-14 13-14	15 15 15
Ciprofloxacin (5mcg) cip			

BIOMARK Laboratories-INDIAwww.biomarklabs.com**TECHNICAL SHEET**

EC (ATCC 25922)	15	16-20	21
SA (ATCC 25923)	15	16-20	21
PS (ATCC 27853)	15	16-20	21
Augmentin (30mcg) aug/ amoxyclav			
EC (ATCC 25922)	13	14-17	18
SA (ATCC 25923)	19	--	20
PS (ATCC 27853)	--	--	--
Vancomycin (30mcg) van			
EC (ATCC 25922)	--	--	--
SA (ATCC 25923)	14	15-16	17
PS (ATCC 27853)	--	--	--
Ceftazidime (10mcg) cpz			
EC (ATCC 25922)	--	--	--
SA (ATCC 25923)	--	--	--
PS (ATCC 27853)	--	--	--
Cephalexin (1.5mcg) cp			
EC (ATCC 25922)	--	--	--
SA (ATCC 25923)	--	--	--
PS (ATCC 27853)	--	--	--

USE: It is used for antimicrobial sensitivity for relevant test organisms for laboratory diagnosis.

Precautions :For Laboratory Use.

Storage: Below 8°C (preferably at -20°C)

Packing: 50 rings/100 rings Pack.

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARK LABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.