

TECHNICAL SHEET

BF010

POTASSIUM TELLURITE 1%

FORMULA

Ingredients:

Potassium tellurite 0.1 gm
Distilled water 10.00 ml

Directions:

Warm up the refrigerated vial to room temperature.
Add in appropriate quantity in sterile, molten, cooled (45-50°C) Baird Parker Agar Base(B1142) / Vogel Johnson Agar Base w/o Tellurite (B352) / Mycoplasma Broth Base w/ CV(B266) / TPEY Agar Base(B329) / Tellurite Glycine Agar Base (B770) / Cholera Medium Base (B984) /Dextrose Proteose Peptone Agar Base (B492)/ Cystine Tellurite Agar Base (B479)/ Diphtheria Virulence Agar Base (B500) Tryptone Tellurite Agar Base (B802)/ Baird Staphylococcus Enrichment Broth Base (B415)/ Tellurite Blood Agar Base (B085) / Mitis Salivarius Agar Base (B258) /Monsur Medium Base (B594)/Giolitti-Cantoni Broth (B1434) as desired. Mix well and dispense in sterile Petri plates or tubes.

Precautions :

1. For Laboratory Use.
2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.

Use: Recommended for the selective isolation of Staphylococci and Corynebacteria.

Storage: Between 2 to 8°C.

Packing: 5 vials

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.
The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.