

TECHNICAL SHEET

BF013

CAMPYLOBACTER SUPPLEMENT-I(BLASER-WANG)

FORMULA

Ingredients:

(Per vial, sufficient for 500 ml medium)

Polymyxin	1250.000 IU
Vancomycin	5.000 mg
Trimethoprim	2.500 mg
Amphotericin B	1.000 mg
Cephalothin	7.500 mg

Directions: Rehydrate the contents of 1 vial aseptically with 5 ml sterile distilled water. Mix well to dissolve avoid frothing of the solution. Aseptically add the contents to 500 ml sterile molten cooled (45-50°C) Blood Agar Base no 2 (B116) / Blood Agar Base no 2(w/Agar 1.2%) (B117) / Columbia Blood Agar Base(B994) / Columbia Blood Agar Base (w/Agar 1%) (B148) with 5-7% v/v laked horse blood and rehydrated contents of 1 vial of Campylobacter growth supplement (BF016) / Modified Brucella Agar Base (B690)/Campylobacter Agar Base (B964) /Christopher's Semisolid Brucella Medium Base(B988) /Campylo Thioglycollate Medium base (B1159)/Brucella Broth Base (B126). Mix gently and pour dispense into sterile Petri Plates / tubes.

Precautions :

1. For Laboratory Use.
2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.

Use: Campylobacter supplement recommended for the isolation of campylobacter species

Storage: Between 2 to 8°C.

Packing: 5 vials