

**BF014**

**CAMPYLOBACTER SUPPLEMENT-II (BUTZLER, MODIFIED)**

**FORMULA**

**Ingredients:**

(Per vial, sufficient for 500 ml medium)

Bacitracin	12500.000 IU
Colistin sulphate	5000.000 units
Amphotericin B	5.000 mg
Cephazolin sodium	7.500 mg
Novobiocin	2.500 mg

**Directions:**

Rehydrate the content of one vial aseptically with 5ml of 50% aqueous Ethanol and mix well to dissolve. Avoid frothing of the solution. Aseptically add the contents to 500 ml of sterile molten cooled (45-50°C) Columbia Blood Agar Base (B994) / Columbia Blood Agar Base w/1% Agar (B148) / Blood Agar Base N0 2 (B116)/Blood Agar Base N0 2 w/1.2% Agar(B117) /Brucella Broth Base (B126)/For 470 ml of Modified Brucella Agar Base (B690) along with 5%v/v defibrinated sheep blood. Mix gently and pour into sterile petri plates.

**Precautions :**

1. For Laboratory Use.
2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.

**Use:**

Campylobacter supplement recommended for the selective isolation of campylobacter species

**Storage:**

Between 2 to 8°C.

**Packing:**

5 vials