

BF045

TOLUIDINE BLUE (0.1 GM/VIAL)

FORMULA

Ingredients :

(Per vial sufficient for 1000 ml medium)

Toluidine blue 0.1g

Directions :

Rehydrate the contents of 1 vial in 5 ml of distilled water and add in 1000 ml of rehydrated DNase Test Agar Base (B158). Heat to boiling to dissolve the medium completely. Sterilize by autoclaving the medium at 15 lbs pressure (121°C) for 15 minutes. Cool and pour in sterile petri plates. Alternatively 0.1% aqueous solution of toluidine blue can be used to flood the growth on DNase Test Agar plates.

Precautions :

1. For Laboratory Use.
2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.

Use : This supplement is recommended for detection of deoxyribonuclease activity

Storage: Store below 30°C.

Packing : 5 vial

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.