

BF078	ROSOLIC ACID
FORMULA	
Ingredients:	
Rosolic acid	0.1 g per vial
Directions:	
Dissolve the contents of one vial in 10 ml of 0.2 N NaOH Mix well and aseptically add it to 990 ml M-FC Agar (B871), M-FC Agar Base, Modified (B870), M-FC Broth (B1100). Boil to dissolve the medium completely.	
Precautions :	
1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.	
Use:	Rosolic acid solution is recommended for selective isolation of coliform bacteria.
Storage:	Between 10 to 30°C.
Packing:	5 vials

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARK LABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.