

**BF103**

**CAN SUPPLEMENT**

**FORMULA**

**Ingredients:**

(Per vial, sufficient for 1000 ml medium)

Colistin Sulphate	10.00 mg
Nalidixic acid	10.00 mg

**Directions:**

Rehydrate the contents of 1 vial aseptically with 2 ml of sterile distilled water. Mix well. Aseptically add it to 950 ml of sterile, molten, cooled (45-50°C) Schaedler Agar(B276) along with 50ml sterile defibrinated sheep blood and Vitamin K1 Supplement(BF118). Mix well and pour into sterile petri plates.

**Precautions :**

1. For Laboratory Use.
2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.

**Use:** A selective supplement recommended for isolation of anaerobic gram positive cocci.

**Storage:** Between 2 to 8°C.

**Packing :**5 vial

**Disclaimer:**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.