

BF117I FRASER SELECTIVE SUPPLEMENT

FORMULA

Ingredients:

(per vial sufficient for 1000 ml medium)

Acriflavin hydrochloride	12.50 mg
Nalidixic acid	10.00 mg

Directions:

Rehydrate the contents of 1 vial aseptically with 10 ml of sterile distilled water. Mix well and aseptically add to 990 ml of sterile, cooled (45-50°C) Fraser Broth Base (B963I). Mix well and dispense as desired.

Precautions :

1. For Laboratory Use.
2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.

Use : An antimicrobial supplement recommended for selective isolation and cultivation of *Listeria monocytogenes* from food and environmental specimens.

Storage: Between 2 to 8°C.

Packing : 5 vials

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.