

BI692 Liver Hydrolysate Enzymatic

Specifications

Appearance : Reddish brown, homogeneous, free flowing powder with characteristic taste and odour

Total Nitrogen : Not less than 11.0%
α – Amino nitrogen : Not less than 3.5%
Moisture : Not more than 6.0%
Ash : Not more than 12.5%
Sodium chloride : Not more than 5.4%

Clarity (2% solution) : Clear

pH (2% solution) : 5 – 7

Microbial Load:(CFU/g)

Total aerobic microbial count by plate method when incubated at 30-35 °c for 48 hours

Bacterial count : NMT 5000

Total Yeast and mold count by plate method when incubated at 20-25 °c for 120 hours

Yeast and mold count : NMT 250

Test for specific organisms

E.coli : Absent
Salmonella species : Absent
Pseudomonas aeruginosa : Absent
Staphylococcus aureus : Absent

Application

Owing to its nutritive characteristics, it is recommended for cultivation of fastidious anaerobic bacteria such as Brucella and Clostridia by adding to Thioglycollate media. It is also incorporated in Blood Agar Base No.2 (B116) for cultivation of wide variety of pathogenic microorganisms. For bulk production of vaccine, steroids, enzymes and other products.

Precaution

1. For Laboratory Use
2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.

Storage : Below 30°c in cool, dry place.

Packing : 500 g